

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
1 February 2001 (01.02.2001)

PCT

(10) International Publication Number
WO 01/07628 A2

(51) International Patent Classification⁷: C12N 15/52,
9/00, C07K 16/40, C12Q 1/68, A61K 38/43, 48/00

(21) International Application Number: PCT/US00/19980

(22) International Filing Date: 20 July 2000 (20.07.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/144,992 22 July 1999 (22.07.1999) US
60/168,858 2 December 1999 (02.12.1999) US

(63) Related by continuation (CON) or continuation-in-part
(CIP) to earlier applications:

US 60/168,858 (CIP)

Filed on 2 December 1999 (02.12.1999)

US 60/144,992 (CIP)

Filed on 22 July 1999 (22.07.1999)

(71) Applicant (for all designated States except US): INCYTE
GENOMICS, INC. [US/US]; 3160 Porter Drive, Palo
Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): TANG, Y., Tom
[CN/US]; 4230 Ranwick Court, San Jose, CA 95118 (US).
HILLMAN, Jennifer, L. [US/US]; 230 Monroe Drive,
#12, Mountain View, CA 94040 (US). BANDMAN, Olga
[US/US]; 366 Anna Avenue, Mountain View, CA 94043
(US). YUE, Henry [US/US]; 826 Lois Avenue, Sunnyvale,

CA 94087 (US). BAUGHN, Mariah, R. [US/US]; 14244
Santiago Road, San Leandro, CA 94577 (US). LAL,
Preeti [IN/US]; 2382 Lass Drive, Santa Clara, CA 95054
(US). LU, Dyung, Aina, M. [US/US]; 55 Park Belmont
Place, San Jose, CA 95136 (US). SHAH, Purvi [IN/US];
1608 Queen Charlotte Drive, #5, Sunnyvale, CA 94087
(US). AZIMZAI, Yalda [US/US]; 2045 Rock Springs
Drive, Hayward, CA 94545 (US).

(74) Agents: HAMLET-COX, Diana et al.; Incyte Genomics,
Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).

(81) Designated States (national): AE, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK,
DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL,
IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,
LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT,
RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA,
UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— Without international search report and to be republished
upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: HUMAN SYNTHETASES

(57) Abstract: The invention provides human synthetases (SYNT) and polynucleotides which identify and encode SYNT. The in-
vention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for
diagnosing, treating, or preventing disorders associated with expression of SYNT.



WO 01/07628 A2

HUMAN SYNTHETASES

5

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of synthetases and to the use of these sequences in the diagnosis, treatment, and prevention of immune, neuronal, and reproductive disorders, and cell proliferative disorders including cancer.

10

BACKGROUND OF THE INVENTION

A large number of cellular biosynthetic intermediary metabolism processes involve intermolecular transfer of carbon atom-containing substrates (carbon substrates). Examples of such reactions include the tricarboxylic acid cycle, synthesis of fatty acids and long-chain phospholipids, synthesis of alcohols and aldehydes, synthesis of intermediary metabolites, and reactions involved in the amino acid degradation pathways. Many of these reactions are catalyzed by synthetases (also called ligases), which catalyze the formation of a bond between two substrate molecules. Some of these reactions require input of energy, usually in the form of conversion of ATP to either ADP or AMP and pyrophosphate. Synthetases are named for the products of the reaction they catalyze and are involved in such processes as metabolism and the synthesis of macromolecules.

20 Ligases forming carbon-oxygen bonds

Proteins make up more than half of the total dry mass of a cell. The synthesis of proteins is central to cell maintenance, growth, and development. Synthesis occurs on ribosomes and depends on the cooperative interaction of several classes of RNA molecules. The process begins with transcription of the genetic code contained within the DNA to form messenger RNA (mRNA). The mRNA moves in steps through a ribosome and the nucleotide sequence of the mRNA is translated into a corresponding sequence of amino acids to construct a distinct protein chain.

The aminoacyl-transfer RNA (tRNA) synthetases are important RNA-associated enzymes with roles in translation. Protein biosynthesis depends on each amino acid forming a linkage with the appropriate tRNA. The aminoacyl-tRNA synthetases are responsible for the activation and correct attachment of an amino acid with its cognate tRNA. The 20 aminoacyl-tRNA synthetase enzymes can be divided into two structural classes, and each class is characterized by a distinctive topology of the catalytic domain. Class I enzymes contain a catalytic domain based on the nucleotide-binding Rossman 'fold'. Class II enzymes contain a central catalytic domain, which consists of a seven-stranded antiparallel β -sheet motif, as well as N- and C- terminal regulatory domains. Class II enzymes are separated into two groups based on the heterodimeric or homodimeric structure of the enzyme; the latter group is further subdivided by the structure of the N- and C-terminal regulatory

domains (Hartlein, M. and Cusack, S. (1995) *J. Mol. Evol.* 40:519-530). Autoantibodies against aminoacyl-tRNAs are generated by patients with dermatomyositis and polymyositis, and correlate strongly with complicating interstitial lung disease (ILD). These antibodies appear to be generated in response to viral infection, and coxsackie virus has been used to induce experimental viral myositis in animals (Friedman, A.W. et al. (1996) *Semin. Arthritis Rheum.* 26:459-467). A synthetase homolog has been shown to be expressed in chronic myeloid leukemia (CML). A phenylalanine-tRNA synthetase homolog has been found to be tumor-selective and expressed in a cell cycle stage- and differentiation-dependent fashion in an acute-phase human CML cell line (Sen, S. et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:6164-6169).

10 Ligases forming carbon-sulfur bonds (Acid-thiol ligases)

In many cases, a carbon substrate is derived from a small molecule containing at least two carbon atoms. The carbon substrate is often covalently bound to a larger molecule which acts as a carbon substrate carrier molecule within the cell. In the biosynthetic mechanisms described above, the carrier molecule is coenzyme A. Coenzyme A (CoA) is structurally related to derivatives of the nucleotide ADP and consists of 4'-phosphopantetheine linked via a phosphodiester bond to the alpha phosphate group of adenosine 3',5'-bisphosphate. The terminal thiol group of 4'-phosphopantetheine acts as the site for carbon substrate bond formation. The predominant carbon substrates which utilize CoA as a carrier molecule during biosynthesis and intermediary metabolism in the cell are acetyl, succinyl, and propionyl moieties, collectively referred to as acyl groups. Other carbon substrates include enoyl lipid, which acts as a fatty acid oxidation intermediate, and carnitine, which acts as an acetyl-CoA flux regulator/ mitochondrial acyl group transfer protein. Acyl-CoA and acetyl-CoA are synthesized in the cell by acyl-CoA synthetase and acetyl-CoA synthetase, respectively.

Activation of fatty acids is mediated by at least three forms of acyl-CoA synthetase activity:

i) acetyl-CoA synthetase, which activates acetate and several other low molecular weight carboxylic acids and is found in muscle mitochondria and the cytosol of other tissues; ii) medium-chain acyl-CoA synthetase, which activates fatty acids containing between four and eleven carbon atoms (predominantly from dietary sources), and is present only in liver mitochondria; and iii) acyl CoA synthetase, which is specific for long chain fatty acids with between six and twenty carbon atoms, and is found in microsomes and the mitochondria. Proteins associated with acyl-CoA synthetase activity have been identified from many sources including bacteria, yeast, plants, mouse, and man. The activity of acyl-CoA synthetase may be modulated by phosphorylation of the enzyme by cAMP-dependent protein kinase. The COL4A5 (collagen, type IV, alpha-5) chromosomal region found deleted in 2 patients with Alport syndrome, elliptocytosis, and mental retardation (Piccini, M. et al. (1998) *Genomics* 47: 350-358) is contiguous with the region containing long-chain acyl-CoA synthetase 4 (FACL4). Therefore, it has been suggested (Piccini supra) that the absence of FACL4 may be involved in the development of mental retardation and other phenotypes associated with

Alport syndrome in these patients.

Ligases forming carbon-nitrogen bonds

A key representative of the amide synthases is the enzyme glutamine synthetase (glutamate-ammonia ligase) that catalyzes the amination of glutamic acid to glutamine by ammonia using the energy of ATP hydrolysis. Glutamine is the primary source for the amino group in various amide transfer reactions involved in de novo pyrimidine nucleotide synthesis and in purine and pyrimidine ribonucleotide interconversions, as well as the conversion of aspartate to asparagine. Overexpression of glutamine synthetase has been observed in primary liver cancer (Christa, L. et al. (1994) Gastroent. 106:1312-1320).

Cyclo-ligases and other carbon-nitrogen ligases comprise various enzymes and enzyme complexes that participate in the de novo pathways to purine and pyrimidine biosynthesis. Because these pathways are critical to the synthesis of nucleotides for replication of both RNA and DNA, many of these enzymes have been the targets of clinical agents for the treatment of cell proliferative disorders such as cancer and infectious diseases.

Purine biosynthesis occurs de novo from the amino acids glycine and glutamine, and other small molecules. Three of the key reactions in this process are catalyzed by a trifunctional enzyme composed of glycinamide-ribonucleotide synthetase (GARS), aminoimidazole ribonucleotide synthetase (AIRS), and glycinamide ribonucleotide transformylase (GART). Together these three enzymes combine ribosylamine phosphate with glycine to yield phosphoribosyl aminoimidazole, a precursor to both adenylylate and guanylate nucleotides. This trifunctional protein has been implicated in the pathology of Downs syndrome (Aimi, J. et al. (1990) Nucleic Acid Res. 18:6665-6672). Adenylosuccinate synthetase catalyzes a later step in purine biosynthesis that converts inosinic acid to adenylosuccinate, a key step on the path to ATP synthesis. This enzyme is also similar to another carbon-nitrogen ligase, argininosuccinate synthetase, that catalyzes a similar reaction in the urea cycle (Powell, S.M. et al. (1992) FEBS Lett. 303:4-10).

Like the de novo biosynthesis of purines, de novo synthesis of the pyrimidine nucleotides uridylylate and cytidylylate also arises from a common precursor, in this instance the nucleotide orotidylylate derived from orotate and phosphoribosyl pyrophosphate (PPRP). Again a trifunctional enzyme comprising three carbon-nitrogen ligases plays a key role in the process. In this case the enzymes aspartate transcarbamylase (ATCase), carbamyl phosphate synthetase II, and dihydroorotase (DHOase) are encoded by a single gene called CAD. Together these three enzymes combine the initial reactants in pyrimidine biosynthesis, glutamine, CO₂ and ATP to form dihydroorotate, the precursor to orotate and orotidylylate (Iwahana, H. et al. (1996) Biochem. Biophys. Res. Commun. 219:249-255). Further steps then lead to the synthesis of uridine nucleotides from orotidylylate. Cytidine nucleotides are derived from uridine-5'-triphosphate (UTP) by the amidation of UTP using glutamine as the amino donor and the enzyme CTP synthetase. Regulatory mutations in the human

CTP synthetase are believed to confer multi-drug resistance to agents widely used in cancer therapy (Yamauchi, M. et al. (1990) EMBO J. 9:2095-2099).

Ligases forming carbon-carbon bonds

Ligases in this group are represented by the carboxylases acetyl-CoA carboxylase and pyruvate carboxylase. Acetyl-CoA carboxylase is a complex which includes a biotin carboxyl carrier protein, biotin carboxylase, and a carboxyl transferase made up of two alpha and two beta subunits. This complex catalyzes the carboxylation of Acetyl-CoA from CO₂ and H₂O using the energy of ATP hydrolysis (PRINTS document PR01069). Acetyl-CoA carboxylase is the rate-limiting step in the biogenesis of long-chain fatty acids. Two isoforms of Acetyl-CoA carboxylase, types I and types II, are expressed in humans in a tissue-specific manner (Ha, J. et al. (1994) Eur. J. Biochem. 219:297-306). Pyruvate carboxylase is a nuclear-encoded mitochondrial enzyme that catalyzes the conversion of pyruvate to oxaloacetate, a key intermediate in the citric acid cycle.

Ligases forming phosphoric ester bonds

Ligases in this group are represented by the DNA ligases involved in both DNA replication and repair. DNA ligases seal phosphodiester bonds between two adjacent nucleotides in a DNA chain using the energy from ATP hydrolysis to first activate the free 5'-phosphate of one nucleotide and then react it with the 3'-OH group of the adjacent nucleotide. This resealing reaction is used in both DNA replication to join small DNA fragments called "Okazaki" fragments that are transiently formed in the process of replicating new DNA, and in DNA repair. DNA repair is the process by which accidental base changes, such as those produced by oxidative damage, hydrolytic attack, or uncontrolled methylation of DNA, are corrected before replication or transcription of the DNA can occur. Bloom's syndrome is an inherited human disease in which individuals are partially deficient in DNA ligation and consequently have an increased incidence of cancer (Alberts, B. et al. (1994) The Molecular Biology of the Cell, Garland Publishing Inc., New York, NY, p. 247).

The discovery of new synthetases and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of immune, neuronal, and reproductive disorders, and cell proliferative disorders including cancer.

SUMMARY OF THE INVENTION

The invention features purified polypeptides, human synthetases, referred to collectively as "SYNT" and individually as "SYNT-1," "SYNT-2," "SYNT-3," "SYNT-4," "SYNT-5," "SYNT-6," "SYNT-7," "SYNT-8," "SYNT-9," "SYNT-10," "SYNT-11," "SYNT-12," "SYNT-13," "SYNT-14," and "SYNT-15." In one aspect, the invention provides an isolated polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID

NO:1-15, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-15. In one alternative, the invention provides an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:1-15.

5 The invention further provides an isolated polynucleotide encoding a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, c) a biologically active fragment of an amino acid sequence selected from the group
10 consisting of SEQ ID NO:1-15, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-15. In one alternative, the polynucleotide encodes a polypeptide selected from the group consisting of SEQ ID NO:1-15. In another alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:16-30.

 Additionally, the invention provides a recombinant polynucleotide comprising a promoter
15 sequence operably linked to a polynucleotide encoding a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, c) a biologically active fragment of an amino acid sequence selected from the group consisting of
20 SEQ ID NO:1-15, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-15. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

 The invention also provides a method for producing a polypeptide comprising an amino acid
25 sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, and d) an immunogenic fragment of an amino acid sequence selected from the
30 group consisting of SEQ ID NO:1-15. The method comprises a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

 Additionally, the invention provides an isolated antibody which specifically binds to a
35 polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, b) a naturally occurring amino

acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-15.

5 The invention further provides an isolated polynucleotide comprising a polynucleotide sequence selected from the group consisting of a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:16-30, b) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:16-30, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence
10 complementary to b), and e) an RNA equivalent of a)-d). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

 Additionally, the invention provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide comprising a polynucleotide sequence selected from the group consisting of a) a polynucleotide sequence selected from the group
15 consisting of SEQ ID NO:16-30, b) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:16-30, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). The method comprises a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence
20 complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and b) detecting the presence or absence of said hybridization complex, and optionally, if present, the amount thereof. In one alternative, the probe comprises at least 60 contiguous nucleotides.

25 The invention further provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide comprising a polynucleotide sequence selected from the group consisting of a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:16-30, b) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID
30 NO:16-30, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). The method comprises a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

35 The invention further provides a pharmaceutical composition comprising an effective amount of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an

amino acid sequence selected from the group consisting of SEQ ID NO:1-15, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, and a pharmaceutically acceptable excipient. In one embodiment, the pharmaceutical composition comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1-15. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional SYNT, comprising administering to a patient in need of such treatment the pharmaceutical composition.

The invention also provides a method for screening a compound for effectiveness as an agonist of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-15. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample. In one alternative, the invention provides a pharmaceutical composition comprising an agonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with decreased expression of functional SYNT, comprising administering to a patient in need of such treatment the pharmaceutical composition.

Additionally, the invention provides a method for screening a compound for effectiveness as an antagonist of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-15. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting antagonist activity in the sample. In one alternative, the invention provides a pharmaceutical composition comprising an antagonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with overexpression of functional SYNT, comprising administering to a patient in need of such treatment the pharmaceutical composition.

The invention further provides a method of screening for a compound that specifically binds to a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-15. The method comprises a) combining the polypeptide with at least one test compound under suitable conditions, and b) detecting binding of the polypeptide to the test compound, thereby identifying a compound that specifically binds to the polypeptide.

The invention further provides a method of screening for a compound that modulates the activity of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-15, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-15. The method comprises a) combining the polypeptide with at least one test compound under conditions permissive for the activity of the polypeptide, b) assessing the activity of the polypeptide in the presence of the test compound, and c) comparing the activity of the polypeptide in the presence of the test compound with the activity of the polypeptide in the absence of the test compound, wherein a change in the activity of the polypeptide in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide.

The invention further provides a method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence selected from the group consisting of SEQ ID NO:16-30, the method comprising a) exposing a sample comprising the target polynucleotide to a compound, and b) detecting altered expression of the target polynucleotide.

The invention further provides a method for assessing toxicity of a test compound, said method comprising a) treating a biological sample containing nucleic acids with the test compound; b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide comprising a polynucleotide sequence selected from the group consisting of i) a polynucleotide sequence selected from the group consisting of SEQ ID NO:16-30, ii) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:16-30, iii) a polynucleotide sequence complementary to i), iv) a polynucleotide sequence complementary to ii),

and v) an RNA equivalent of i)-iv). Hybridization occurs under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:16-30, ii) a naturally occurring polynucleotide sequence having at least
5 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:16-30, iii) a polynucleotide sequence complementary to i), iv) a polynucleotide sequence complementary to ii), and v) an RNA equivalent of i)-iv). Alternatively, the target polynucleotide comprises a fragment of the above polynucleotide sequence; c) quantifying the amount of hybridization complex; and d) comparing the amount of hybridization complex in the treated
10 biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

BRIEF DESCRIPTION OF THE TABLES

15 Table 1 shows polypeptide and nucleotide sequence identification numbers (SEQ ID NOs), clone identification numbers (clone IDs), cDNA libraries, and cDNA fragments used to assemble full-length sequences encoding SYNT.

Table 2 shows features of each polypeptide sequence, including potential motifs, homologous sequences, and methods, algorithms, and searchable databases used for analysis of SYNT.

20 Table 3 shows selected fragments of each nucleic acid sequence; the tissue-specific expression patterns of each nucleic acid sequence as determined by northern analysis; diseases, disorders, or conditions associated with these tissues; and the vector into which each cDNA was cloned.

Table 4 describes the tissues used to construct the cDNA libraries from which cDNA clones
25 encoding SYNT were isolated.

Table 5 shows the tools, programs, and algorithms used to analyze the polynucleotides and polypeptides of the invention, along with applicable descriptions, references, and threshold parameters.

DESCRIPTION OF THE INVENTION

30 Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which
35 will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an,"

and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

5 Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing
10 the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

DEFINITIONS

"SYNT" refers to the amino acid sequences of substantially purified SYNT obtained from
15 any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist" refers to a molecule which intensifies or mimics the biological activity of SYNT. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of SYNT either by directly interacting with
20 SYNT or by acting on components of the biological pathway in which SYNT participates.

An "allelic variant" is an alternative form of the gene encoding SYNT. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to
25 allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

"Altered" nucleic acid sequences encoding SYNT include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as SYNT or a
30 polypeptide with at least one functional characteristic of SYNT. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding SYNT, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding SYNT. The encoded protein may also be "altered," and may contain deletions, insertions, or
35 substitutions of amino acid residues which produce a silent change and result in a functionally equivalent SYNT. Deliberate amino acid substitutions may be made on the basis of similarity in

polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of SYNT is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

The terms "amino acid" and "amino acid sequence" refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where "amino acid sequence" is recited to refer to a sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

"Amplification" relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term "antagonist" refers to a molecule which inhibits or attenuates the biological activity of SYNT. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of SYNT either by directly interacting with SYNT or by acting on components of the biological pathway in which SYNT participates.

The term "antibody" refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')₂, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind SYNT polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition capable of base-pairing with the "sense"

(coding) strand of a specific nucleic acid sequence. Antisense compositions may include DNA; RNA; peptide nucleic acid (PNA); oligonucleotides having modified backbone linkages such as phosphorothioates, methylphosphonates, or benzylphosphonates; oligonucleotides having modified sugar groups such as 2'-methoxyethyl sugars or 2'-methoxyethoxy sugars; or oligonucleotides having modified bases such as 5-methyl cytosine, 2'-deoxyuracil, or 7-deaza-2'-deoxyguanosine. Antisense molecules may be produced by any method including chemical synthesis or transcription. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand of a reference DNA molecule.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" or "immunogenic" refers to the capability of the natural, recombinant, or synthetic SYNT, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

"Complementary" describes the relationship between two single-stranded nucleic acid sequences that anneal by base-pairing. For example, 5'-AGT-3' pairs with its complement, 3'-TCA-5'.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding SYNT or fragments of SYNT may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been subjected to repeated DNA sequence analysis to resolve uncalled bases, extended using the XL-PCR kit (PE Biosystems, Foster City CA) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from one or more overlapping cDNA, EST, or genomic DNA fragments using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI) or Phrap (University of Washington, Seattle WA). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that are predicted to least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows

amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

	Original Residue	Conservative Substitution
	Ala	Gly, Ser
5	Arg	His, Lys
	Asn	Asp, Gln, His
	Asp	Asn, Glu
	Cys	Ala, Ser
	Gln	Asn, Glu, His
10	Glu	Asp, Gln, His
	Gly	Ala
	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
	Leu	Ile, Val
15	Lys	Arg, Gln, Glu
	Met	Leu, Ile
	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
	Thr	Ser, Val
20	Trp	Phe, Tyr
	Tyr	His, Phe, Trp
	Val	Ile, Leu, Thr

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to a chemically modified polynucleotide or polypeptide. Chemical modifications of a polynucleotide sequence can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

A "detectable label" refers to a reporter molecule or enzyme that is capable of generating a measurable signal and is covalently or noncovalently joined to a polynucleotide or polypeptide.

A "fragment" is a unique portion of SYNT or the polynucleotide encoding SYNT which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10,

15, 16, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50% of a polypeptide) as shown in a certain
5 defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

A fragment of SEQ ID NO:16-30 comprises a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:16-30, for example, as distinct from any other sequence in the
10 genome from which the fragment was obtained. A fragment of SEQ ID NO:16-30 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:16-30 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:16-30 and the region of SEQ ID NO:16-30 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

15 A fragment of SEQ ID NO:1-15 is encoded by a fragment of SEQ ID NO:16-30. A fragment of SEQ ID NO:1-15 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-15. For example, a fragment of SEQ ID NO:1-15 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-15. The precise length of a fragment of SEQ ID NO:1-15 and the region of SEQ ID NO:1-15 to which the fragment
20 corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A "full-length" polynucleotide sequence is one containing at least a translation initiation codon (e.g., methionine) followed by an open reading frame and a translation termination codon. A "full-length" polynucleotide sequence encodes a "full-length" polypeptide sequence.

25 "Homology" refers to sequence similarity or, interchangeably, sequence identity, between two or more polynucleotide sequences or two or more polypeptide sequences.

The terms "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps
30 in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of
35 molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS

8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polynucleotide sequences.

- 5 Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence
- 10 analysis programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST
- 15 programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Reward for match: 1

20 *Penalty for mismatch: -2*

Open Gap: 5 and Extension Gap: 2 penalties

Gap x drop-off: 50

Expect: 10

Word Size: 11

25 *Filter: on*

- Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous
- 30 nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

- Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes
- 35 in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to the percentage of residue matches between at least two polypeptide sequences aligned using a standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail above, generally preserve the charge and hydrophobicity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default residue weight table. As with polynucleotide alignments, the percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.12 (Apr-21-2000) with blastp set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Open Gap: 11 and Extension Gap: 1 penalties

Gap x drop-off: 50

Expect: 10

Word Size: 3

Filter: on

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

"Human artificial chromosomes" (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for chromosome replication, segregation and maintenance.

The term "humanized antibody" refers to an antibody molecule in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

“Hybridization” refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of complementarity. Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the “washing” step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml sheared, denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Such wash temperatures are typically selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating T_m and conditions for nucleic acid hybridization are well known and can be found in Sambrook, J. et al., 1989, Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, sheared and denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term “hybridization complex” refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C_{ot} or R_{ot} analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate).

to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

5 "Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

10 An "immunogenic fragment" is a polypeptide or oligopeptide fragment of SYNT which is capable of eliciting an immune response when introduced into a living organism, for example, a mammal. The term "immunogenic fragment" also includes any polypeptide or oligopeptide fragment of SYNT which is useful in any of the antibody production methods disclosed herein or known in the art.

The term "microarray" refers to an arrangement of a plurality of polynucleotides, polypeptides, or other chemical compounds on a substrate.

15 The terms "element" and "array element" refer to a polynucleotide, polypeptide, or other chemical compound having a unique and defined position on a microarray.

The term "modulate" refers to a change in the activity of SYNT. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of SYNT.

20 The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

25 "Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with a second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

30 "Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

35 "Post-translational modification" of an SYNT may involve lipidation, glycosylation, phosphorylation, acetylation, racemization, proteolytic cleavage, and other modifications known in the art. These processes may occur synthetically or biochemically. Biochemical modifications will

vary by cell type depending on the enzymatic milieu of SYNT.

"Probe" refers to nucleic acid sequences encoding SYNT, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule.

5 Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes.

"Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

10 Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the
15 specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for example Sambrook, J. et al., 1989, Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel, F.M. et al., 1987, Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis, M. et al., 1990, PCR
20 Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

Oligonucleotides for use as primers are selected using software known in the art for such
25 purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of
30 Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the
35 selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific

needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both
5 unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

10 A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, *supra*. The term recombinant includes nucleic acids that have
15 been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a
20 vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

A "regulatory element" refers to a nucleic acid sequence usually derived from untranslated regions of a gene and includes enhancers, promoters, introns, and 5' and 3' untranslated regions (UTRs). Regulatory elements interact with host or viral proteins which control transcription,
25 translation, or RNA stability.

"Reporter molecules" are chemical or biochemical moieties used for labeling a nucleic acid, amino acid, or antibody. Reporter molecules include radionuclides; enzymes; fluorescent, chemiluminescent, or chromogenic agents; substrates; cofactors; inhibitors; magnetic particles; and other moieties known in the art.

30 An "RNA equivalent," in reference to a DNA sequence, is composed of the same linear sequence of nucleotides as the reference DNA sequence with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The term "sample" is used in its broadest sense. A sample suspected of containing nucleic
35 acids encoding SYNT, or fragments thereof, or SYNT itself, may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or

cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide comprising the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acid residues or nucleotides by different amino acid residues or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

A "transcript image" refers to the collective pattern of gene expression by a particular cell type or tissue under given conditions at a given time.

"Transformation" describes a process by which exogenous DNA is introduced into a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, bacteriophage or viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed" cells includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "transgenic organism," as used herein, is any organism, including but not limited to animals and plants, in which one or more of the cells of the organism contains heterologous nucleic acid introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or in vitro fertilization, but rather is directed to the introduction of a recombinant DNA molecule. The

transgenic organisms contemplated in accordance with the present invention include bacteria, cyanobacteria, fungi, plants, and animals. The isolated DNA of the present invention can be introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention into such organisms are widely known and provided in references such as Sambrook et al. (1989), supra.

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternative splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% or greater sequence identity over a certain defined length of one of the polypeptides.

THE INVENTION

The invention is based on the discovery of new human synthetases (SYNT), the polynucleotides encoding SYNT, and the use of these compositions for the diagnosis, treatment, or prevention of immune, neuronal, and reproductive disorders, and cell proliferative disorders including cancer.

Table 1 lists the Incyte clones used to assemble full length nucleotide sequences encoding SYNT. Columns 1 and 2 show the sequence identification numbers (SEQ ID NOs) of the polypeptide

and nucleotide sequences, respectively. Column 3 shows the clone IDs of the Incyte clones in which nucleic acids encoding each SYNT were identified, and column 4 shows the cDNA libraries from which these clones were isolated. Column 5 shows Incyte clones and their corresponding cDNA libraries. Clones for which cDNA libraries are not indicated were derived from pooled cDNA

5 libraries. In some cases, GenBank sequence identifiers are also shown in column 5. The Incyte clones and GenBank cDNA sequences, where indicated, in column 5 were used to assemble the consensus nucleotide sequence of each SYNT and are useful as fragments in hybridization technologies.

The columns of Table 2 show various properties of each of the polypeptides of the invention: column 1 references the SEQ ID NO; column 2 shows the number of amino acid residues in each
10 polypeptide; column 3 shows potential phosphorylation sites; column 4 shows potential glycosylation sites; column 5 shows the amino acid residues comprising signature sequences and motifs; column 6 shows homologous sequences as identified by BLAST analysis; and column 7 shows analytical methods and in some cases, searchable databases to which the analytical methods were applied. The methods of column 7 were used to characterize each polypeptide through sequence homology and
15 protein motifs.

The columns of Table 3 show the tissue-specificity and diseases, disorders, or conditions associated with nucleotide sequences encoding SYNT. The first column of Table 3 lists the nucleotide SEQ ID NOs. Column 2 lists fragments of the nucleotide sequences of column 1. These fragments are useful, for example, in hybridization or amplification technologies to identify SEQ ID
20 NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19 SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:29, and SEQ ID NO:30 and to distinguish between SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19 SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:29, and SEQ ID NO:30 and related polynucleotide sequences.
25 The polypeptides encoded by these fragments are useful, for example, as immunogenic peptides. Column 3 lists tissue categories which express SYNT as a fraction of total tissues expressing SYNT. Column 4 lists diseases, disorders, or conditions associated with those tissues expressing SYNT as a fraction of total tissues expressing SYNT. Column 5 lists the vectors used to subclone each cDNA library.

30 The columns of Table 4 show descriptions of the tissues used to construct the cDNA libraries from which cDNA clones encoding SYNT were isolated. Column 1 references the nucleotide SEQ ID NOs, column 2 shows the cDNA libraries from which these clones were isolated, and column 3 shows the tissue origins and other descriptive information relevant to the cDNA libraries in column 2.

SEQ ID NO: 16 maps to chromosome 5 within the interval from 147.10 to 150.00
35 centiMorgans. SEQ ID NO: 17 maps to chromosome 10 within the interval from 137.60 to 139.20 centiMorgans. This interval also contains gene MX11, a member of the MYC family. SEQ ID NO:

18 maps to chromosome 2 within the interval from 228.80 to 230.10 centiMorgans. This interval also contains a gene for a proto-oncogene encoding a tyrosine-protein kinase. SEQ ID NO:21 maps to chromosome 5 within the interval from 172.6 to 184.7 centiMorgans. SEQ ID NO:24 maps to chromosome 2 within the interval from 118.0 to 127.4 centiMorgans. SEQ ID NO:26 maps to
5 chromosome 3 within the interval from 157.4 to 162.0 centiMorgans. SEQ ID NO:27 maps to chromosome 12 within the interval from 97.1 to 116.6 centiMorgans. SEQ ID NO:28 maps to chromosome 4 within the interval from 77.3 to 99.2 centiMorgans and to chromosome 5 within the intervals from 79.2 to 92.3 centiMorgans, from 116.3 to 127.9 centiMorgans, and from 157.6 to 163.0 centiMorgans. SEQ ID NO:29 maps to chromosome 1 within the interval from 242.5 to 258.7
10 centiMorgans and to chromosome 19 within the interval from 69.9 to 104.9 centiMorgans. SEQ ID NO:30 maps to chromosome 1 within the interval from 57.2 to 57.5 centiMorgans.

The invention also encompasses SYNT variants. A preferred SYNT variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the SYNT amino acid sequence, and which contains at least one functional or
15 structural characteristic of SYNT.

The invention also encompasses polynucleotides which encode SYNT. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:16-30, which encodes SYNT. The polynucleotide sequences of SEQ ID NO:16-30, as presented in the Sequence Listing, embrace the equivalent RNA
20 sequences, wherein occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The invention also encompasses a variant of a polynucleotide sequence encoding SYNT. In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to the polynucleotide
25 sequence encoding SYNT. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:16-30 which has at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:16-30. Any one of the polynucleotide variants described above can encode an amino
30 acid sequence which contains at least one functional or structural characteristic of SYNT.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding SYNT, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide
35 sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the

polynucleotide sequence of naturally occurring SYNT, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode SYNT and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring SYNT under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding SYNT or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding SYNT and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode SYNT and SYNT derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding SYNT or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:16-30 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (PE Biosystems, Foster City CA), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (PE Biosystems). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (PE Biosystems), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit

7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding SYNT may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) PCR Methods Applic. 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) Nucleic Acids Res. 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) PCR Methods Applic. 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) Nucleic Acids Res. 19:3055-3060). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, PE Biosystems), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments

which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode SYNT may be cloned in recombinant DNA molecules that direct expression of SYNT, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent
5 degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express SYNT.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter SYNT-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA
10 shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

The nucleotides of the present invention may be subjected to DNA shuffling techniques such
15 as MOLECULARBREEDING (Maxygen Inc., Santa Clara CA; described in U.S. Patent Number 5,837,458; Chang, C.-C. et al. (1999) Nat. Biotechnol. 17:793-797; Christians, F.C. et al. (1999) Nat. Biotechnol. 17:259-264; and Cramer, A. et al. (1996) Nat. Biotechnol. 14:315-319) to alter or improve the biological properties of SYNT, such as its biological or enzymatic activity or its ability to bind to other molecules or compounds. DNA shuffling is a process by which a library of gene
20 variants is produced using PCR-mediated recombination of gene fragments. The library is then subjected to selection or screening procedures that identify those gene variants with the desired properties. These preferred variants may then be pooled and further subjected to recursive rounds of DNA shuffling and selection/screening. Thus, genetic diversity is created through "artificial" breeding and rapid molecular evolution. For example, fragments of a single gene containing random
25 point mutations may be recombined, screened, and then reshuffled until the desired properties are optimized. Alternatively, fragments of a given gene may be recombined with fragments of homologous genes in the same gene family, either from the same or different species, thereby maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable manner.

In another embodiment, sequences encoding SYNT may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) Nucleic Acids Symp. Ser. 7:215-223; and Horn, T. et al. (1980) Nucleic Acids Symp. Ser. 7:225-232.)
Alternatively, SYNT itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solution-phase or solid-phase techniques.
35 (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY, pp. 55-60; and Roberge, J.Y. et al. (1995) Science 269:202-204.) Automated synthesis

may be achieved using the ABI 431A peptide synthesizer (PE Biosystems). Additionally, the amino acid sequence of SYNT, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide or a polypeptide having a sequence of a naturally occurring polypeptide.

5 The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) *Methods Enzymol.* 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, *supra*, pp. 28-53.)

10 In order to express a biologically active SYNT, the nucleotide sequences encoding SYNT or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding SYNT. Such elements may vary in their strength and specificity. Specific initiation signals
15 may also be used to achieve more efficient translation of sequences encoding SYNT. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding SYNT and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted,
20 exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

25 Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding SYNT and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et
30 al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

35 A variety of expression vector/host systems may be utilized to contain and express sequences encoding SYNT. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV,

- or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. (See, e.g., Sambrook, supra; Ausubel, supra; Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509; Bitter, G.A. et al. (1987) Methods Enzymol. 153:516-544; Scorer, C.A. et al. (1994) Bio/Technology 12:181-184; Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945; Takamatsu, N. (1987) EMBO J. 6:307-311; Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al. (1984) Science 224:838-843; Winter, J. et al. (1991) Results Probl. Cell Differ. 17:85-105; The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196; Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659; and Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.) Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. (See, e.g., Di Nicola, M. et al. (1998) Cancer Gen. Ther. 5(6):350-356; Yu, M. et al., (1993) Proc. Natl. Acad. Sci. USA 90(13):6340-6344; Buller, R.M. et al. (1985) Nature 317(6040):813-815; McGregor, D.P. et al. (1994) Mol. Immunol. 31(3):219-226; and Verma, I.M. and N. Somia (1997) Nature 389:239-242.)
- The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding SYNT. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding SYNT can be achieved using a multifunctional *E. coli* vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORT1 plasmid (Life Technologies). Ligation of sequences encoding SYNT into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for *in vitro* transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of SYNT are needed, e.g. for the production of antibodies, vectors which direct high level expression of SYNT may be used. For example, vectors containing the strong, inducible T5 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of SYNT. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast *Saccharomyces cerevisiae* or *Pichia pastoris*. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, supra; Bitter, supra; and Scorer, supra.)

Plant systems may also be used for expression of SYNT. Transcription of sequences encoding SYNT may be driven viral promoters, e.g., the 35S and 19S promoters of CaMV used alone

or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J. 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, *supra*; Broglie, *supra*; and Winter, *supra*.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding SYNT may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses SYNT in host cells. (See, e.g., Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of SYNT in cell lines is preferred. For example, sequences encoding SYNT can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk* and *apv* cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981)

J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech), β glucuronidase and its substrate β -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding SYNT is inserted within a marker gene sequence, transformed cells containing sequences encoding SYNT can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding SYNT under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding SYNT and that express SYNT may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of SYNT using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on SYNT is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding SYNT include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding SYNT, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase

such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding SYNT may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode SYNT may be designed to contain signal sequences which direct secretion of SYNT through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding SYNT may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric SYNT protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of SYNT activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the SYNT encoding sequence and the heterologous protein sequence, so that SYNT may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, supra, ch. 10).

A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled SYNT may be achieved in vitro using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These
5 systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, ³⁵S-methionine.

SYNT of the present invention or fragments thereof may be used to screen for compounds that specifically bind to SYNT. At least one and up to a plurality of test compounds may be screened
10 for specific binding to SYNT. Examples of test compounds include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

In one embodiment, the compound thus identified is closely related to the natural ligand of SYNT, e.g., a ligand or fragment thereof, a natural substrate, a structural or functional mimetic, or a natural binding partner. (See, Coligan, J.E. et al. (1991) Current Protocols in Immunology 1(2):
15 Chapter 5.) Similarly, the compound can be closely related to the natural receptor to which SYNT binds, or to at least a fragment of the receptor, e.g., the ligand binding site. In either case, the compound can be rationally designed using known techniques. In one embodiment, screening for these compounds involves producing appropriate cells which express SYNT, either as a secreted protein or on the cell membrane. Preferred cells include cells from mammals, yeast, Drosophila, or
20 E. coli. Cells expressing SYNT or cell membrane fractions which contain SYNT are then contacted with a test compound and binding, stimulation, or inhibition of activity of either SYNT or the compound is analyzed.

An assay may simply test binding of a test compound to the polypeptide, wherein binding is detected by a fluorophore, radioisotope, enzyme conjugate, or other detectable label. For example,
25 the assay may comprise the steps of combining at least one test compound with SYNT, either in solution or affixed to a solid support, and detecting the binding of SYNT to the compound. Alternatively, the assay may detect or measure binding of a test compound in the presence of a labeled competitor. Additionally, the assay may be carried out using cell-free preparations, chemical libraries, or natural product mixtures, and the test compound(s) may be free in solution or affixed to a
30 solid support.

SYNT of the present invention or fragments thereof may be used to screen for compounds that modulate the activity of SYNT. Such compounds may include agonists, antagonists, or partial or inverse agonists. In one embodiment, an assay is performed under conditions permissive for SYNT activity, wherein SYNT is combined with at least one test compound, and the activity of SYNT in the
35 presence of a test compound is compared with the activity of SYNT in the absence of the test compound. A change in the activity of SYNT in the presence of the test compound is indicative of a

compound that modulates the activity of SYNT. Alternatively, a test compound is combined with an in vitro or cell-free system comprising SYNT under conditions suitable for SYNT activity, and the assay is performed. In either of these assays, a test compound which modulates the activity of SYNT may do so indirectly and need not come in direct contact with the test compound. At least one and up to a plurality of test compounds may be screened.

In another embodiment, polynucleotides encoding SYNT or their mammalian homologs may be "knocked out" in an animal model system using homologous recombination in embryonic stem (ES) cells. Such techniques are well known in the art and are useful for the generation of animal models of human disease. (See, e.g., U.S. Patent No. 5,175,383 and U.S. Patent No. 5,767,337.) For example, mouse ES cells, such as the mouse 129/SvJ cell line, are derived from the early mouse embryo and grown in culture. The ES cells are transformed with a vector containing the gene of interest disrupted by a marker gene, e.g., the neomycin phosphotransferase gene (neo; Capecchi, M.R. (1989) Science 244:1288-1292). The vector integrates into the corresponding region of the host genome by homologous recombination. Alternatively, homologous recombination takes place using the Cre-loxP system to knockout a gene of interest in a tissue- or developmental stage-specific manner (Marth, J.D. (1996) Clin. Invest. 97:1999-2002; Wagner, K.U. et al. (1997) Nucleic Acids Res. 25:4323-4330). Transformed ES cells are identified and microinjected into mouse cell blastocysts such as those from the C57BL/6 mouse strain. The blastocysts are surgically transferred to pseudopregnant dams, and the resulting chimeric progeny are genotyped and bred to produce heterozygous or homozygous strains. Transgenic animals thus generated may be tested with potential therapeutic or toxic agents.

Polynucleotides encoding SYNT may also be manipulated in vitro in ES cells derived from human blastocysts. Human ES cells have the potential to differentiate into at least eight separate cell lineages including endoderm, mesoderm, and ectodermal cell types. These cell lineages differentiate into, for example, neural cells, hematopoietic lineages, and cardiomyocytes (Thomson, J.A. et al. (1998) Science 282:1145-1147).

Polynucleotides encoding SYNT can also be used to create "knockin" humanized animals (pigs) or transgenic animals (mice or rats) to model human disease. With knockin technology, a region of a polynucleotide encoding SYNT is injected into animal ES cells, and the injected sequence integrates into the animal cell genome. Transformed cells are injected into blastulae, and the blastulae are implanted as described above. Transgenic progeny or inbred lines are studied and treated with potential pharmaceutical agents to obtain information on treatment of a human disease. Alternatively, a mammal inbred to overexpress SYNT, e.g., by secreting SYNT in its milk, may also serve as a convenient source of that protein (Janne, J. et al. (1998) Biotechnol. Annu. Rev. 4:55-74).

35 THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists

between regions of SYNT and human synthetases. In addition, the expression of SYNT is closely associated with hematopoietic/immune, cancerous, proliferating, inflamed, immune, nervous, gastrointestinal and reproductive tissues. Therefore, SYNT appears to play a role in an immune disorder such as inflammation, actinic keratosis, acquired immunodeficiency syndrome (AIDS),

5 Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, arteriosclerosis, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, bronchitis, bursitis, cholecystitis, cirrhosis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease,

10 Hashimoto's thyroiditis, paroxysmal nocturnal hemoglobinuria, hepatitis, hypereosinophilia, irritable bowel syndrome, episodic lymphopenia with lymphocytotoxins, mixed connective tissue disease (MCTD), multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, myelofibrosis, osteoarthritis, osteoporosis, pancreatitis, polycythemia vera, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis,

15 systemic lupus erythematosus, systemic sclerosis, primary thrombocythemia, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, trauma, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and hematopoietic cancer including lymphoma, leukemia, and myeloma; a neuronal disorder, such as akathisia, Alzheimer's disease, amnesia, amyotrophic lateral sclerosis, bipolar

20 disorder, catatonia, cerebral neoplasms, dementia, depression, diabetic neuropathy, Down's syndrome, tardive dyskinesia, dystonias, epilepsy, Huntington's disease, peripheral neuropathy, multiple sclerosis, neurofibromatosis, Parkinson's disease, paranoid psychoses, postherpetic neuralgia, schizophrenia, and Tourette's disorder; a reproductive disorder, such as a disorder of prolactin production, infertility, including tubal disease, ovulatory defects, and endometriosis, a disruption of

25 the estrous cycle, a disruption of the menstrual cycle, polycystic ovary syndrome, ovarian hyperstimulation syndrome, an endometrial or ovarian tumor, a uterine fibroid, autoimmune disorders, an ectopic pregnancy, and teratogenesis; cancer of the breast, fibrocystic breast disease, and galactorrhea; a disruption of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence,

30 carcinoma of the male breast, and gynecomastia; and a cell proliferative disorder, such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, a cancer of the adrenal gland, bladder, bone,

35 bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis,

thymus, thyroid, and uterus.

In another embodiment, a vector capable of expressing SYNT or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of SYNT including, but not limited to, those described above.

5 In a further embodiment, a pharmaceutical composition comprising a substantially purified SYNT in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of SYNT including, but not limited to, those provided above.

10 In still another embodiment, an agonist which modulates the activity of SYNT may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of SYNT including, but not limited to, those listed above.

In a further embodiment, an antagonist of SYNT may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of SYNT. Examples of such disorders include, but are not limited to, those immune, neuronal, reproductive, and cell proliferative
15 disorders described above. In one aspect, an antibody which specifically binds SYNT may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express SYNT.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding SYNT may be administered to a subject to treat or prevent a disorder associated with
20 increased expression or activity of SYNT including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The
25 combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of SYNT may be produced using methods which are generally known in the art. In particular, purified SYNT may be used to produce antibodies or to screen libraries of
30 pharmaceutical agents to identify those which specifically bind SYNT. Antibodies to SYNT may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

35 For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with SYNT or with any fragment or oligopeptide thereof

which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to SYNT have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein. Short stretches of SYNT amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to SYNT may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) *Nature* 256:495-497; Kozbor, D. et al. (1985) *J. Immunol. Methods* 81:31-42; Cote, R.J. et al. (1983) *Proc. Natl. Acad. Sci. USA* 80:2026-2030; and Cole, S.P. et al. (1984) *Mol. Cell Biol.* 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) *Proc. Natl. Acad. Sci. USA* 81:6851-6855; Neuberger, M.S. et al. (1984) *Nature* 312:604-608; and Takeda, S. et al. (1985) *Nature* 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce SYNT-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) *Proc. Natl. Acad. Sci. USA* 88:10134-10137.)

Antibodies may also be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:3833-3837; Winter, G. et al. (1991) *Nature* 349:293-299.)

Antibody fragments which contain specific binding sites for SYNT may also be generated. For example, such fragments include, but are not limited to, F(ab')₂ fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) *Science* 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between SYNT and its
5 specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering SYNT epitopes is generally used, but a competitive binding assay may also be employed (Pound, supra).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for SYNT. Affinity is expressed as an
10 association constant, K_a , which is defined as the molar concentration of SYNT-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The K_a determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple SYNT epitopes, represents the average affinity, or avidity, of the antibodies for SYNT. The K_a determined for a preparation of monoclonal antibodies, which are monospecific for a
15 particular SYNT epitope, represents a true measure of affinity. High-affinity antibody preparations with K_a ranging from about 10^9 to 10^{12} L/mole are preferred for use in immunoassays in which the SYNT-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K_a ranging from about 10^6 to 10^7 L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of SYNT, preferably in active form, from the
20 antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington DC; Liddell, J.E. and A. Cryer (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For
25 example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of SYNT-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al., supra.)

30 In another embodiment of the invention, the polynucleotides encoding SYNT, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, modifications of gene expression can be achieved by designing complementary sequences or antisense molecules (DNA, RNA, PNA, or modified oligonucleotides) to the coding or regulatory regions of the gene encoding SYNT. Such technology is well known in the art, and antisense oligonucleotides or larger
35 fragments can be designed from various locations along the coding or control regions of sequences encoding SYNT. (See, e.g., Agrawal, S., ed. (1996) Antisense Therapeutics, Humana Press Inc.,

Totawa NJ.)

In therapeutic use, any gene delivery system suitable for introduction of the antisense sequences into appropriate target cells can be used. Antisense sequences can be delivered intracellularly in the form of an expression plasmid which, upon transcription, produces a sequence complementary to at least a portion of the cellular sequence encoding the target protein. (See, e.g., Slater, J.E. et al. (1998) *J. Allergy Clin. Immunol.* 102(3):469-475; and Scanlon, K.J. et al. (1995) 9(13):1288-1296.) Antisense sequences can also be introduced intracellularly through the use of viral vectors, such as retrovirus and adeno-associated virus vectors. (See, e.g., Miller, A.D. (1990) *Blood* 76:271; Ausubel, *supra*; Uckert, W. and W. Walther (1994) *Pharmacol. Ther.* 63(3):323-347.) Other gene delivery mechanisms include liposome-derived systems, artificial viral envelopes, and other systems known in the art. (See, e.g., Rossi, J.J. (1995) *Br. Med. Bull.* 51(1):217-225; Boado, R.J. et al. (1998) *J. Pharm. Sci.* 87(11):1308-1315; and Morris, M.C. et al. (1997) *Nucleic Acids Res.* 25(14):2730-2736.)

In another embodiment of the invention, polynucleotides encoding SYNT may be used for somatic or germline gene therapy. Gene therapy may be performed to (i) correct a genetic deficiency (e.g., in the cases of severe combined immunodeficiency (SCID)-X1 disease characterized by X-linked inheritance (Cavazzana-Calvo, M. et al. (2000) *Science* 288:669-672), severe combined immunodeficiency syndrome associated with an inherited adenosine deaminase (ADA) deficiency (Blaese, R.M. et al. (1995) *Science* 270:475-480; Bordignon, C. et al. (1995) *Science* 270:470-475), cystic fibrosis (Zabner, J. et al. (1993) *Cell* 75:207-216; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:643-666; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:667-703), thalassemias, familial hypercholesterolemia, and hemophilia resulting from Factor VIII or Factor IX deficiencies (Crystal, R.G. (1995) *Science* 270:404-410; Verma, I.M. and Somia, N. (1997) *Nature* 389:239-242)), (ii) express a conditionally lethal gene product (e.g., in the case of cancers which result from unregulated cell proliferation), or (iii) express a protein which affords protection against intracellular parasites (e.g., against human retroviruses, such as human immunodeficiency virus (HIV) (Baltimore, D. (1988) *Nature* 335:395-396; Poeschla, E. et al. (1996) *Proc. Natl. Acad. Sci. USA.* 93:11395-11399), hepatitis B or C virus (HBV, HCV); fungal parasites, such as Candida albicans and Paracoccidioides brasiliensis; and protozoan parasites such as Plasmodium falciparum and Trypanosoma cruzi). In the case where a genetic deficiency in SYNT expression or regulation causes disease, the expression of SYNT from an appropriate population of transduced cells may alleviate the clinical manifestations caused by the genetic deficiency.

In a further embodiment of the invention, diseases or disorders caused by deficiencies in SYNT are treated by constructing mammalian expression vectors encoding SYNT and introducing these vectors by mechanical means into SYNT-deficient cells. Mechanical transfer technologies for use with cells in vivo or ex vitro include (i) direct DNA microinjection into individual cells, (ii)

ballistic gold particle delivery, (iii) liposome-mediated transfection, (iv) receptor-mediated gene transfer, and (v) the use of DNA transposons (Morgan, R.A. and W.F. Anderson (1993) *Annu. Rev. Biochem.* 62:191-217; Ivics, Z. (1997) *Cell* 91:501-510; Boulay, J.-L. and H. R  cipon (1998) *Curr. Opin. Biotechnol.* 9:445-450).

5 Expression vectors that may be effective for the expression of SYNT include, but are not limited to, the pCDNA 3.1, EPITAG, PRCCMV2, PREP, PVAX vectors (Invitrogen, Carlsbad CA), PCMV-SCRIPT, PCMV-TAG, PEGSH/PERV (Stratagene, La Jolla CA), and PTET-OFF, PTET-ON, PTRE2, PTRE2-LUC, PTK-HYG (Clontech, Palo Alto CA). SYNT may be expressed using (i) a constitutively active promoter, (e.g., from cytomegalovirus (CMV), Rous sarcoma virus
10 (RSV), SV40 virus, thymidine kinase (TK), or β -actin genes), (ii) an inducible promoter (e.g., the tetracycline-regulated promoter (Gossen, M. and H. Bujard (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551; Gossen, M. et al. (1995) *Science* 268:1766-1769; Rossi, F.M.V. and H.M. Blau (1998) *Curr. Opin. Biotechnol.* 9:451-456), commercially available in the T-REX plasmid (Invitrogen)); the ecdysone-inducible promoter (available in the plasmids PVGRXR and PIND; Invitrogen); the
15 FK506/rapamycin inducible promoter; or the RU486/mifepristone inducible promoter (Rossi, F.M.V. and H.M. Blau, *supra*)), or (iii) a tissue-specific promoter or the native promoter of the endogenous gene encoding SYNT from a normal individual.

 Commercially available liposome transformation kits (e.g., the PERFECT LIPID TRANSFECTION KIT, available from Invitrogen) allow one with ordinary skill in the art to deliver
20 polynucleotides to target cells in culture and require minimal effort to optimize experimental parameters. In the alternative, transformation is performed using the calcium phosphate method (Graham, F.L. and A.J. Eb (1973) *Virology* 52:456-467), or by electroporation (Neumann, E. et al. (1982) *EMBO J.* 1:841-845). The introduction of DNA to primary cells requires modification of these standardized mammalian transfection protocols.

25 In another embodiment of the invention, diseases or disorders caused by genetic defects with respect to SYNT expression are treated by constructing a retrovirus vector consisting of (i) the polynucleotide encoding SYNT under the control of an independent promoter or the retrovirus long terminal repeat (LTR) promoter, (ii) appropriate RNA packaging signals, and (iii) a Rev-responsive element (RRE) along with additional retrovirus *cis*-acting RNA sequences and coding sequences
30 required for efficient vector propagation. Retrovirus vectors (e.g., PFB and PFBNEO) are commercially available (Stratagene) and are based on published data (Riviere, I. et al. (1995) *Proc. Natl. Acad. Sci. USA* 92:6733-6737), incorporated by reference herein. The vector is propagated in an appropriate vector producing cell line (VPCL) that expresses an envelope gene with a tropism for receptors on the target cells or a promiscuous envelope protein such as VSVg (Armentano, D. et al.
35 (1987) *J. Virol.* 61:1647-1650; Bender, M.A. et al. (1987) *J. Virol.* 61:1639-1646; Adam, M.A. and A.D. Miller (1988) *J. Virol.* 62:3802-3806; Dull, T. et al. (1998) *J. Virol.* 72:8463-8471; Zufferey, R.

et al. (1998) J. Virol. 72:9873-9880). U.S. Patent Number 5,910,434 to Rigg ("Method for obtaining retrovirus packaging cell lines producing high transducing efficiency retroviral supernatant") discloses a method for obtaining retrovirus packaging cell lines and is hereby incorporated by reference. Propagation of retrovirus vectors, transduction of a population of cells (e.g., CD4⁺ T-cells), and the return of transduced cells to a patient are procedures well known to persons skilled in the art of gene therapy and have been well documented (Ranga, U. et al. (1997) J. Virol. 71:7020-7029; Bauer, G. et al. (1997) Blood 89:2259-2267; Bonyhadi, M.L. (1997) J. Virol. 71:4707-4716; Ranga, U. et al. (1998) Proc. Natl. Acad. Sci. USA 95:1201-1206; Su, L. (1997) Blood 89:2283-2290).

In the alternative, an adenovirus-based gene therapy delivery system is used to deliver polynucleotides encoding SYNT to cells which have one or more genetic abnormalities with respect to the expression of SYNT. The construction and packaging of adenovirus-based vectors are well known to those with ordinary skill in the art. Replication defective adenovirus vectors have proven to be versatile for importing genes encoding immunoregulatory proteins into intact islets in the pancreas (Csete, M.E. et al. (1995) Transplantation 27:263-268). Potentially useful adenoviral vectors are described in U.S. Patent Number 5,707,618 to Armentano ("Adenovirus vectors for gene therapy"), hereby incorporated by reference. For adenoviral vectors, see also Antinozzi, P.A. et al. (1999) Annu. Rev. Nutr. 19:511-544; and Verma, I.M. and N. Somia (1997) Nature 18:389:239-242, both incorporated by reference herein.

In another alternative, a herpes-based, gene therapy delivery system is used to deliver polynucleotides encoding SYNT to target cells which have one or more genetic abnormalities with respect to the expression of SYNT. The use of herpes simplex virus (HSV)-based vectors may be especially valuable for introducing SYNT to cells of the central nervous system, for which HSV has a tropism. The construction and packaging of herpes-based vectors are well known to those with ordinary skill in the art. A replication-competent herpes simplex virus (HSV) type 1-based vector has been used to deliver a reporter gene to the eyes of primates (Liu, X. et al. (1999) Exp. Eye Res. 169:385-395). The construction of a HSV-1 virus vector has also been disclosed in detail in U.S. Patent Number 5,804,413 to DeLuca ("Herpes simplex virus strains for gene transfer"), which is hereby incorporated by reference. U.S. Patent Number 5,804,413 teaches the use of recombinant HSV d92 which consists of a genome containing at least one exogenous gene to be transferred to a cell under the control of the appropriate promoter for purposes including human gene therapy. Also taught by this patent are the construction and use of recombinant HSV strains deleted for ICP4, ICP27 and ICP22. For HSV vectors, see also Goins, W.F. et al. (1999) J. Virol. 73:519-532 and Xu, H. et al. (1994) Dev. Biol. 163:152-161, hereby incorporated by reference. The manipulation of cloned herpesvirus sequences, the generation of recombinant virus following the transfection of multiple plasmids containing different segments of the large herpesvirus genomes, the growth and propagation

of herpesvirus, and the infection of cells with herpesvirus are techniques well known to those of ordinary skill in the art.

In another alternative, an alphavirus (positive, single-stranded RNA virus) vector is used to deliver polynucleotides encoding SYNT to target cells. The biology of the prototypic alphavirus, Semliki Forest Virus (SFV), has been studied extensively and gene transfer vectors have been based on the SFV genome (Garoff, H. and K.-J. Li (1998) Curr. Opin. Biotech. 9:464-469). During alphavirus RNA replication, a subgenomic RNA is generated that normally encodes the viral capsid proteins. This subgenomic RNA replicates to higher levels than the full-length genomic RNA, resulting in the overproduction of capsid proteins relative to the viral proteins with enzymatic activity (e.g., protease and polymerase). Similarly, inserting the coding sequence for SYNT into the alphavirus genome in place of the capsid-coding region results in the production of a large number of SYNT-coding RNAs and the synthesis of high levels of SYNT in vector transduced cells. While alphavirus infection is typically associated with cell lysis within a few days, the ability to establish a persistent infection in hamster normal kidney cells (BHK-21) with a variant of Sindbis virus (SIN) indicates that the lytic replication of alphaviruses can be altered to suit the needs of the gene therapy application (Dryga, S.A. et al. (1997) Virology 228:74-83). The wide host range of alphaviruses will allow the introduction of SYNT into a variety of cell types. The specific transduction of a subset of cells in a population may require the sorting of cells prior to transduction. The methods of manipulating infectious cDNA clones of alphaviruses, performing alphavirus cDNA and RNA transfections, and performing alphavirus infections, are well known to those with ordinary skill in the art.

Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may also be employed to inhibit gene expression. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding SYNT.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by

scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding SYNT. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

An additional embodiment of the invention encompasses a method for screening for a compound which is effective in altering expression of a polynucleotide encoding SYNT. Compounds which may be effective in altering expression of a specific polynucleotide may include, but are not limited to, oligonucleotides, antisense oligonucleotides, triple helix-forming oligonucleotides, transcription factors and other polypeptide transcriptional regulators, and non-macromolecular chemical entities which are capable of interacting with specific polynucleotide sequences. Effective compounds may alter polynucleotide expression by acting as either inhibitors or promoters of polynucleotide expression. Thus, in the treatment of disorders associated with increased SYNT expression or activity, a compound which specifically inhibits expression of the polynucleotide encoding SYNT may be therapeutically useful, and in the treatment of disorders associated with decreased SYNT expression or activity, a compound which specifically promotes expression of the polynucleotide encoding SYNT may be therapeutically useful.

At least one, and up to a plurality, of test compounds may be screened for effectiveness in altering expression of a specific polynucleotide. A test compound may be obtained by any method

commonly known in the art, including chemical modification of a compound known to be effective in altering polynucleotide expression; selection from an existing, commercially-available or proprietary library of naturally-occurring or non-natural chemical compounds; rational design of a compound based on chemical and/or structural properties of the target polynucleotide; and selection from a

5 library of chemical compounds created combinatorially or randomly. A sample comprising a polynucleotide encoding SYNT is exposed to at least one test compound thus obtained. The sample may comprise, for example, an intact or permeabilized cell, or an *in vitro* cell-free or reconstituted biochemical system. Alterations in the expression of a polynucleotide encoding SYNT are assayed by any method commonly known in the art. Typically, the expression of a specific nucleotide is
10 detected by hybridization with a probe having a nucleotide sequence complementary to the sequence of the polynucleotide encoding SYNT. The amount of hybridization may be quantified, thus forming the basis for a comparison of the expression of the polynucleotide both with and without exposure to one or more test compounds. Detection of a change in the expression of a polynucleotide exposed to a test compound indicates that the test compound is effective in altering the expression of the
15 polynucleotide. A screen for a compound effective in altering expression of a specific polynucleotide can be carried out, for example, using a *Schizosaccharomyces pombe* gene expression system (Atkins, D. et al. (1999) U.S. Patent No. 5,932,435; Arndt, G.M. et al. (2000) Nucleic Acids Res. 28:E15) or a human cell line such as HeLa cell (Clarke, M.L. et al. (2000) Biochem. Biophys. Res. Commun. 268:8-13). A particular embodiment of the present invention involves screening a
20 combinatorial library of oligonucleotides (such as deoxyribonucleotides, ribonucleotides, peptide nucleic acids, and modified oligonucleotides) for antisense activity against a specific polynucleotide sequence (Bruce, T.W. et al. (1997) U.S. Patent No. 5,686,242; Bruce, T.W. et al. (2000) U.S. Patent No. 6,022,691).

Many methods for introducing vectors into cells or tissues are available and equally suitable
25 for use *in vivo*, *in vitro*, and *ex vivo*. For *ex vivo* therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

30 Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a pharmaceutical composition which generally comprises an active ingredient formulated with a pharmaceutically
35 acceptable excipient. Excipients may include, for example, sugars, starches, celluloses, gums, and proteins. Various formulations are commonly known and are thoroughly discussed in the latest

edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA). Such pharmaceutical compositions may consist of SYNT, antibodies to SYNT, and mimetics, agonists, antagonists, or inhibitors of SYNT.

The pharmaceutical compositions utilized in this invention may be administered by any
5 number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, pulmonary, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

Pharmaceutical compositions for pulmonary administration may be prepared in liquid or dry powder form. These compositions are generally aerosolized immediately prior to inhalation by the
10 patient. In the case of small molecules (e.g. traditional low molecular weight organic drugs), aerosol delivery of fast-acting formulations is well-known in the art. In the case of macromolecules (e.g. larger peptides and proteins), recent developments in the field of pulmonary delivery via the alveolar region of the lung have enabled the practical delivery of drugs such as insulin to blood circulation (see, e.g., Patton, J.S. et al., U.S. Patent No. 5,997,848). Pulmonary delivery has the advantage of
15 administration without needle injection, and obviates the need for potentially toxic penetration enhancers.

Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

20 Specialized forms of pharmaceutical compositions may be prepared for direct intracellular delivery of macromolecules comprising SYNT or fragments thereof. For example, liposome preparations containing a cell-impermeable macromolecule may promote cell fusion and intracellular delivery of the macromolecule. Alternatively, SYNT or a fragment thereof may be joined to a short cationic N-terminal portion from the HIV Tat-1 protein. Fusion proteins thus generated have been
25 found to transduce into the cells of all tissues, including the brain, in a mouse model system (Schwarze, S.R. et al. (1999) *Science* 285:1569-1572).

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, monkeys, or pigs. An animal model may also be used to determine the appropriate concentration
30 range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example SYNT or fragments thereof, antibodies of SYNT, and agonists, antagonists or inhibitors of SYNT, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by
35 standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the ED₅₀ (the dose therapeutically effective in 50% of the population) or LD₅₀ (the dose

lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the LD_{50}/ED_{50} ratio. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1 μg to 100,000 μg , up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

DIAGNOSTICS

In another embodiment, antibodies which specifically bind SYNT may be used for the diagnosis of disorders characterized by expression of SYNT, or in assays to monitor patients being treated with SYNT or agonists, antagonists, or inhibitors of SYNT. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for SYNT include methods which utilize the antibody and a label to detect SYNT in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring SYNT, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of SYNT expression. Normal or standard values for SYNT expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibody to SYNT under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of SYNT expressed in

subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding SYNT may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of SYNT may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of SYNT, and to monitor regulation of SYNT levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding SYNT or closely related molecules may be used to identify nucleic acid sequences which encode SYNT. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding SYNT, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and may have at least 50% sequence identity to any of the SYNT encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:16-30 or from genomic sequences including promoters, enhancers, and introns of the SYNT gene.

Means for producing specific hybridization probes for DNAs encoding SYNT include the cloning of polynucleotide sequences encoding SYNT or SYNT derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as ³²P or ³⁵S, or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding SYNT may be used for the diagnosis of disorders associated with expression of SYNT. Examples of such disorders include, but are not limited to, an immune disorder such as inflammation, actinic keratosis, acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, arteriosclerosis, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, bronchitis, bursitis, cholecystitis, cirrhosis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, paroxysmal nocturnal hemoglobinuria, hepatitis, hypereosinophilia, irritable bowel syndrome, episodic lymphopenia with lymphocytotoxins, mixed connective tissue

disease (MCTD), multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, myelofibrosis, osteoarthritis, osteoporosis, pancreatitis, polycythemia vera, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, primary thrombocythemia, thrombocytopenic

5 purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, trauma, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and hematopoietic cancer including lymphoma, leukemia, and myeloma; a neuronal disorder, such as akathisia, Alzheimer's disease, amnesia, amyotrophic lateral sclerosis, bipolar disorder, catatonia, cerebral neoplasms, dementia, depression, diabetic neuropathy, Down's syndrome,

10 tardive dyskinesia, dystonias, epilepsy, Huntington's disease, peripheral neuropathy, multiple sclerosis, neurofibromatosis, Parkinson's disease, paranoid psychoses, postherpetic neuralgia, schizophrenia, and Tourette's disorder; a reproductive disorder, such as a disorder of prolactin production, infertility, including tubal disease, ovulatory defects, and endometriosis, a disruption of the estrous cycle, a disruption of the menstrual cycle, polycystic ovary syndrome, ovarian

15 hyperstimulation syndrome, an endometrial or ovarian tumor, a uterine fibroid, autoimmune disorders, an ectopic pregnancy, and teratogenesis; cancer of the breast, fibrocystic breast disease, and galactorrhea; a disruption of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence, carcinoma of the male breast, and gynecomastia; and a cell proliferative disorder, such as actinic

20 keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, a cancer of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver,

25 lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus. The polynucleotide sequences encoding SYNT may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered SYNT expression. Such qualitative or quantitative methods are well known

30 in the art.

In a particular aspect, the nucleotide sequences encoding SYNT may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding SYNT may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a

35 suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to

a control sample then the presence of altered levels of nucleotide sequences encoding SYNT in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

5 In order to provide a basis for the diagnosis of a disorder associated with expression of SYNT, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding SYNT, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from
10 normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated,
15 hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or
20 overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

25 Additional diagnostic uses for oligonucleotides designed from the sequences encoding SYNT may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a polynucleotide encoding SYNT, or a fragment of a polynucleotide complementary to the polynucleotide encoding SYNT, and will be employed under optimized conditions for identification of a specific gene or
30 condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

In a particular aspect, oligonucleotide primers derived from the polynucleotide sequences encoding SYNT may be used to detect single nucleotide polymorphisms (SNPs). SNPs are substitutions, insertions and deletions that are a frequent cause of inherited or acquired genetic
35 disease in humans. Methods of SNP detection include, but are not limited to, single-stranded conformation polymorphism (SSCP) and fluorescent SSCP (fSSCP) methods. In SSCP,

oligonucleotide primers derived from the polynucleotide sequences encoding SYNT are used to amplify DNA using the polymerase chain reaction (PCR). The DNA may be derived, for example, from diseased or normal tissue, biopsy samples, bodily fluids, and the like. SNPs in the DNA cause differences in the secondary and tertiary structures of PCR products in single-stranded form, and these differences are detectable using gel electrophoresis in non-denaturing gels. In fSCCP, the oligonucleotide primers are fluorescently labeled, which allows detection of the amplimers in high-throughput equipment such as DNA sequencing machines. Additionally, sequence database analysis methods, termed in silico SNP (isSNP), are capable of identifying polymorphisms by comparing the sequence of individual overlapping DNA fragments which assemble into a common consensus sequence. These computer-based methods filter out sequence variations due to laboratory preparation of DNA and sequencing errors using statistical models and automated analyses of DNA sequence chromatograms. In the alternative, SNPs may be detected and characterized by mass spectrometry using, for example, the high throughput MASSARRAY system (Sequenom, Inc., San Diego CA).

Methods which may also be used to quantify the expression of SYNT include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) J. Immunol. Methods 159:235-244; Duplaa, C. et al. (1993) Anal. Biochem. 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer or polynucleotide of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as elements on a microarray. The microarray can be used in transcript imaging techniques which monitor the relative expression levels of large numbers of genes simultaneously as described in Seilhamer, J.J. et al., "Comparative Gene Transcript Analysis," U.S. Patent No. 5,840,484, incorporated herein by reference. The microarray may also be used to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, to monitor progression/regression of disease as a function of gene expression, and to develop and monitor the activities of therapeutic agents in the treatment of disease. In particular, this information may be used to develop a pharmacogenomic profile of a patient in order to select the most appropriate and effective treatment regimen for that patient. For example, therapeutic agents which are highly effective and display the fewest side effects may be selected for a patient based on his/her pharmacogenomic profile.

In another embodiment, antibodies specific for SYNT, or SYNT or fragments thereof may be used as elements on a microarray. The microarray may be used to monitor or measure protein-protein interactions, drug-target interactions, and gene expression profiles, as described above.

A particular embodiment relates to the use of the polynucleotides of the present invention to generate a transcript image of a tissue or cell type. A transcript image represents the global pattern of gene expression by a particular tissue or cell type. Global gene expression patterns are analyzed by quantifying the number of expressed genes and their relative abundance under given conditions and at a given time. (See Seilhamer et al., "Comparative Gene Transcript Analysis," U.S. Patent Number 5,840,484, expressly incorporated by reference herein.) Thus a transcript image may be generated by hybridizing the polynucleotides of the present invention or their complements to the totality of transcripts or reverse transcripts of a particular tissue or cell type. In one embodiment, the hybridization takes place in high-throughput format, wherein the polynucleotides of the present invention or their complements comprise a subset of a plurality of elements on a microarray. The resultant transcript image would provide a profile of gene activity.

Transcript images may be generated using transcripts isolated from tissues, cell lines, biopsies, or other biological samples. The transcript image may thus reflect gene expression in vivo, as in the case of a tissue or biopsy sample, or in vitro, as in the case of a cell line.

Transcript images which profile the expression of the polynucleotides of the present invention may also be used in conjunction with in vitro model systems and preclinical evaluation of pharmaceuticals, as well as toxicological testing of industrial and naturally-occurring environmental compounds. All compounds induce characteristic gene expression patterns, frequently termed molecular fingerprints or toxicant signatures, which are indicative of mechanisms of action and toxicity (Nuwaysir, E.F. et al. (1999) Mol. Carcinog. 24:153-159; Steiner, S. and N.L. Anderson (2000) Toxicol. Lett. 112-113:467-471, expressly incorporated by reference herein). If a test compound has a signature similar to that of a compound with known toxicity, it is likely to share those toxic properties. These fingerprints or signatures are most useful and refined when they contain expression information from a large number of genes and gene families. Ideally, a genome-wide measurement of expression provides the highest quality signature. Even genes whose expression is not altered by any tested compounds are important as well, as the levels of expression of these genes are used to normalize the rest of the expression data. The normalization procedure is useful for comparison of expression data after treatment with different compounds. While the assignment of gene function to elements of a toxicant signature aids in interpretation of toxicity mechanisms, knowledge of gene function is not necessary for the statistical matching of signatures which leads to prediction of toxicity. (See, for example, Press Release 00-02 from the National Institute of Environmental Health Sciences, released February 29, 2000, available at <http://www.niehs.nih.gov/oc/news/toxchip.htm>.) Therefore, it is important and desirable in toxicological screening using toxicant signatures to include all expressed gene sequences.

In one embodiment, the toxicity of a test compound is assessed by treating a biological sample containing nucleic acids with the test compound. Nucleic acids that are expressed in the

treated biological sample are hybridized with one or more probes specific to the polynucleotides of the present invention, so that transcript levels corresponding to the polynucleotides of the present invention may be quantified. The transcript levels in the treated biological sample are compared with levels in an untreated biological sample. Differences in the transcript levels between the two samples are indicative of a toxic response caused by the test compound in the treated sample.

Another particular embodiment relates to the use of the polypeptide sequences of the present invention to analyze the proteome of a tissue or cell type. The term proteome refers to the global pattern of protein expression in a particular tissue or cell type. Each protein component of a proteome can be subjected individually to further analysis. Proteome expression patterns, or profiles, are analyzed by quantifying the number of expressed proteins and their relative abundance under given conditions and at a given time. A profile of a cell's proteome may thus be generated by separating and analyzing the polypeptides of a particular tissue or cell type. In one embodiment, the separation is achieved using two-dimensional gel electrophoresis, in which proteins from a sample are separated by isoelectric focusing in the first dimension, and then according to molecular weight by sodium dodecyl sulfate slab gel electrophoresis in the second dimension (Steiner and Anderson, supra). The proteins are visualized in the gel as discrete and uniquely positioned spots, typically by staining the gel with an agent such as Coomassie Blue or silver or fluorescent stains. The optical density of each protein spot is generally proportional to the level of the protein in the sample. The optical densities of equivalently positioned protein spots from different samples, for example, from biological samples either treated or untreated with a test compound or therapeutic agent, are compared to identify any changes in protein spot density related to the treatment. The proteins in the spots are partially sequenced using, for example, standard methods employing chemical or enzymatic cleavage followed by mass spectrometry. The identity of the protein in a spot may be determined by comparing its partial sequence, preferably of at least 5 contiguous amino acid residues, to the polypeptide sequences of the present invention. In some cases, further sequence data may be obtained for definitive protein identification.

A proteomic profile may also be generated using antibodies specific for SYNT to quantify the levels of SYNT expression. In one embodiment, the antibodies are used as elements on a microarray, and protein expression levels are quantified by exposing the microarray to the sample and detecting the levels of protein bound to each array element (Lueking, A. et al. (1999) *Anal. Biochem.* 270:103-111; Mendoze, L.G. et al. (1999) *Biotechniques* 27:778-788). Detection may be performed by a variety of methods known in the art, for example, by reacting the proteins in the sample with a thiol- or amino-reactive fluorescent compound and detecting the amount of fluorescence bound at each array element.

Toxicant signatures at the proteome level are also useful for toxicological screening, and should be analyzed in parallel with toxicant signatures at the transcript level. There is a poor

correlation between transcript and protein abundances for some proteins in some tissues (Anderson, N.L. and J. Seilhamer (1997) *Electrophoresis* 18:533-537), so proteome toxicant signatures may be useful in the analysis of compounds which do not significantly affect the transcript image, but which alter the proteomic profile. In addition, the analysis of transcripts in body fluids is difficult, due to rapid degradation of mRNA, so proteomic profiling may be more reliable and informative in such cases.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins that are expressed in the treated biological sample are separated so that the amount of each protein can be quantified. The amount of each protein is compared to the amount of the corresponding protein in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample. Individual proteins are identified by sequencing the amino acid residues of the individual proteins and comparing these partial sequences to the polypeptides of the present invention.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins from the biological sample are incubated with antibodies specific to the polypeptides of the present invention. The amount of protein recognized by the antibodies is quantified. The amount of protein in the treated biological sample is compared with the amount in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) *Proc. Natl. Acad. Sci. USA* 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.) Various types of microarrays are well known and thoroughly described in DNA Microarrays: A Practical Approach, M. Schena, ed. (1999) Oxford University Press, London, hereby expressly incorporated by reference.

In another embodiment of the invention, nucleic acid sequences encoding SYNT may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. Either coding or noncoding sequences may be used, and in some instances, noncoding sequences may be preferable over coding sequences. For example, conservation of a coding sequence among members of a multi-gene family may potentially cause undesired cross hybridization during chromosomal mapping. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1

constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355; Price, C.M. (1993) *Blood Rev.* 7:127-134; and Trask, B.J. (1991) *Trends Genet.* 7:149-154.) Once mapped, the nucleic acid sequences of the invention may be used to develop genetic linkage maps, for example, which correlate the inheritance of a disease state with the inheritance of a particular chromosome region or restriction fragment length polymorphism (RFLP). (See, e.g., Lander, E.S. and D. Botstein (1986) *Proc. Natl. Acad. Sci. USA* 83:7353-7357.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding SYNT on a physical map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder and thus may further positional cloning efforts.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps.

Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the exact chromosomal locus is not known. This information is valuable to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the gene or genes responsible for a disease or syndrome have been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) *Nature* 336:577-580.) The nucleotide sequence of the instant invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, SYNT, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between SYNT and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with SYNT, or fragments thereof, and washed. Bound SYNT is then detected by methods well known in the art. Purified SYNT can also be coated directly onto plates for use in the aforementioned drug screening techniques.

Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding SYNT specifically compete with a test compound for binding SYNT. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with SYNT.

5 In additional embodiments, the nucleotide sequences which encode SYNT may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding
10 description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications, and publications mentioned above and below, in particular U.S. Ser. No. 60/144,992 and U.S. Ser. No. 60/168,858 are hereby expressly incorporated
15 by reference.

EXAMPLES

I. Construction of cDNA Libraries

RNA was purchased from Clontech or isolated from tissues described in Table 4. Some
20 tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

25 Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A+) RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA
30 purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, supra, units
35 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the

appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g.,

5 PBLUESCRIPT plasmid (Stratagene), PSPORT1 plasmid (Life Technologies), pcDNA2.1 plasmid (Invitrogen, Carlsbad CA), or pINCY plasmid (Incyte Genomics, Palo Alto CA). Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 α , DH10B, or ElectroMAX DH10B from Life Technologies.

II. Isolation of cDNA Clones

10 Plasmids obtained as described in Example I were recovered from host cells by in vivo excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96

15 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in

20 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

III. Sequencing and Analysis

Incyte cDNA recovered in plasmids as described in Example II were sequenced as follows.

25 Sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (PE Biosystems) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI

30 PRISM BIGDYE Terminator cycle sequencing ready reaction kit (PE Biosystems). Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing system (PE Biosystems) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the

35 cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, supra, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in

Example VI.

The polynucleotide sequences derived from cDNA sequencing were assembled and analyzed using a combination of software programs which utilize algorithms well known to those skilled in the art. Table 5 summarizes the tools, programs, and algorithms used and provides applicable descriptions, references, and threshold parameters. The first column of Table 5 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score, the greater the homology between two sequences). Sequences were analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments were generated using the default parameters specified by the clustal algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

The polynucleotide sequences were validated by removing vector, linker, and polyA sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The sequences were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and PFAM to acquire annotation using programs based on BLAST, FASTA, and BLIMPS. The sequences were assembled into full length polynucleotide sequences using programs based on Phred, Phrap, and Consed, and were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length amino acid sequences, and these full length sequences were subsequently analyzed by querying against databases such as the GenBank databases (described above), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and Hidden Markov Model (HMM)-based protein family databases such as PFAM. HMM is a probabilistic approach which analyzes consensus primary structures of gene families. (See, e.g., Eddy, S.R. (1996) Curr. Opin. Struct. Biol. 6:361-365.)

The programs described above for the assembly and analysis of full length polynucleotide and amino acid sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:16-30. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies were described in The Invention section above.

IV. Analysis of Polynucleotide Expression

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs

from a particular cell type or tissue have been bound. (See, e.g., Sambrook, *supra*, ch. 7; Ausubel, 1995, *supra*, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in cDNA databases such as GenBank or LIFESEQ (Incyte Genomics). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\text{BLAST Score} \times \text{Percent Identity}}{5 \times \text{minimum \{length(Seq. 1), length(Seq. 2)\}}}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. The product score is a normalized value between 0 and 100, and is calculated as follows: the BLAST score is multiplied by the percent nucleotide identity and the product is divided by (5 times the length of the shorter of the two sequences). The BLAST score is calculated by assigning a score of +5 for every base that matches in a high-scoring segment pair (HSP), and -4 for every mismatch. Two sequences may share more than one HSP (separated by gaps). If there is more than one HSP, then the pair with the highest BLAST score is used to calculate the product score. The product score represents a balance between fractional overlap and quality in a BLAST alignment. For example, a product score of 100 is produced only for 100% identity over the entire length of the shorter of the two sequences being compared. A product score of 70 is produced either by 100% identity and 70% overlap at one end, or by 88% identity and 100% overlap at the other. A product score of 50 is produced either by 100% identity and 50% overlap at one end, or 79% identity and 100% overlap.

The results of northern analyses are reported as a percentage distribution of libraries in which the transcript encoding SYNT occurred. Analysis involved the categorization of cDNA libraries by organ/tissue and disease. The organ/tissue categories included cardiovascular, dermatologic, developmental, endocrine, gastrointestinal, hematopoietic/immune, musculoskeletal, nervous, reproductive, and urologic. The disease/condition categories included cancer, inflammation, trauma, cell proliferation, neurological, and pooled. For each category, the number of libraries expressing the sequence of interest was counted and divided by the total number of libraries across all categories. Percentage values of tissue-specific and disease- or condition-specific expression are reported in Table 3.

V. Extension of SYNT Encoding Polynucleotides

The full length nucleic acid sequences of SEQ ID NO:16-30 were produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other

primer, to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would
5 result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction
10 mix contained DNA template, 200 nmol of each primer, reaction buffer containing Mg^{2+} , $(NH_4)_2SO_4$, and β -mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the
15 alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100 μ l PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE
20 and 0.5 μ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5 μ l to 10 μ l aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose mini-gel to determine which reactions were successful in extending
25 the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For
30 shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, and individual colonies were picked and cultured overnight at 37°C in
35 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase

(Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (PE Biosystems).

In like manner, the polynucleotide sequences of SEQ ID NO:16-30 are used to obtain 5' regulatory sequences using the procedure above, along with oligonucleotides designed for such extension, and an appropriate genomic library.

V. Chromosomal Mapping of SNYT Encoding Polynucleotides

The cDNA sequences which were used to assemble SEQ ID NO:16-30 were compared with sequences from the Incyte LIFESEQ database and public domain databases using BLAST and other implementations of the Smith-Waterman algorithm. Sequences from these databases that matched SEQ ID NO:16-30 were assembled into clusters of contiguous and overlapping sequences using assembly algorithms such as Phrap (Table 5). Radiation hybrid and genetic mapping data available from public resources such as the Stanford Human Genome Center (SHGC), Whitehead Institute for Genome Research (WIGR), and Généthon were used to determine if any of the clustered sequences had been previously mapped. Inclusion of a mapped sequence in a cluster resulted in the assignment of all sequences of that cluster, including its particular SEQ ID NO., to that map location.

The genetic map locations of SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, and SEQ ID NO:30 are described in The Invention as ranges, or intervals, of human chromosomes. More than one map location is reported for SEQ ID NO:28 and SEQ ID NO:29, indicating that previously mapped sequences having similarity, but not complete identity, to SEQ ID NO:28 and SEQ ID NO:29 were assembled into their respective clusters. The map position of an interval, in centiMorgans, is measured relative to the terminus of the chromosome's p-arm. (The centiMorgan (cM) is a unit of measurement based on recombination frequencies between chromosomal markers. On average, 1 cM is roughly equivalent to 1 megabase (Mb) of DNA in humans, although this can vary widely due to hot and cold spots of recombination.) The cM distances are based on genetic markers mapped by Généthon which provide boundaries for radiation hybrid markers whose sequences were included in each of the clusters. Diseases associated with the public and Incyte sequences located within the indicated intervals are also reported in the Invention where applicable. Human genome maps and other resources available to the public, such as the NCBI "GeneMap'99" World Wide Web site which can be accessed at <http://www.ncbi.nlm.nih.gov/genemap>, can be employed to determine if

previously identified disease genes map within or in proximity to the intervals indicated above.

VI. Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:16-30 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 μ Ci of [γ - 32 P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10^7 counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and compared.

VII. Microarrays

The linkage or synthesis of array elements upon a microarray can be achieved utilizing photolithography, piezoelectric printing (ink-jet printing, See, e.g., Baldeschweiler, *supra*), mechanical microspotting technologies, and derivatives thereof. The substrate in each of the aforementioned technologies should be uniform and solid with a non-porous surface (Schena (1999), *supra*). Suggested substrates include silicon, silica, glass slides, glass chips, and silicon wafers. Alternatively, a procedure analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced using available methods and machines well known to those of ordinary skill in the art and may contain any appropriate number of elements. (See, e.g., Schena, M. et al. (1995) Science 270:467-470; Shalon, D. et al. (1996) Genome Res. 6:639-645; Marshall, A. and J. Hodgson (1998) Nat. Biotechnol. 16:27-31.)

Full length cDNAs, Expressed Sequence Tags (ESTs), or fragments or oligomers thereof may comprise the elements of the microarray. Fragments or oligomers suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). The array elements are hybridized with polynucleotides in a biological sample. The polynucleotides in the biological sample are conjugated to a fluorescent label or other molecular tag for ease of detection.

After hybridization, nonhybridized nucleotides from the biological sample are removed, and a fluorescence scanner is used to detect hybridization at each array element. Alternatively, laser desorption and mass spectrometry may be used for detection of hybridization. The degree of complementarity and the relative abundance of each polynucleotide which hybridizes to an element on the microarray may be assessed. In one embodiment, microarray preparation and usage is described in detail below.

Tissue or Cell Sample Preparation

Total RNA is isolated from tissue samples using the guanidinium thiocyanate method and poly(A)⁺ RNA is purified using the oligo-(dT) cellulose method. Each poly(A)⁺ RNA sample is reverse transcribed using MMLV reverse-transcriptase, 0.05 pg/ μ l oligo-(dT) primer (21mer), 1X first strand buffer, 0.03 units/ μ l RNase inhibitor, 500 μ M dATP, 500 μ M dGTP, 500 μ M dTTP, 40 μ M dCTP, 40 μ M dCTP-Cy3 (BDS) or dCTP-Cy5 (Amersham Pharmacia Biotech). The reverse transcription reaction is performed in a 25 ml volume containing 200 ng poly(A)⁺ RNA with GEMBRIGHT kits (Incyte). Specific control poly(A)⁺ RNAs are synthesized by *in vitro* transcription from non-coding yeast genomic DNA. After incubation at 37 °C for 2 hr, each reaction sample (one with Cy3 and another with Cy5 labeling) is treated with 2.5 ml of 0.5M sodium hydroxide and incubated for 20 minutes at 85 °C to stop the reaction and degrade the RNA. Samples are purified using two successive CHROMA SPIN 30 gel filtration spin columns (CLONTECH Laboratories, Inc. (CLONTECH), Palo Alto CA) and after combining, both reaction samples are ethanol precipitated using 1 ml of glycogen (1 mg/ml), 60 ml sodium acetate, and 300 ml of 100% ethanol. The sample is then dried to completion using a SpeedVAC (Savant Instruments Inc., Holbrook NY) and resuspended in 14 μ l 5X SSC/0.2% SDS.

Microarray Preparation

Sequences of the present invention are used to generate array elements. Each array element is amplified from bacterial cells containing vectors with cloned cDNA inserts. PCR amplification uses primers complementary to the vector sequences flanking the cDNA insert. Array elements are amplified in thirty cycles of PCR from an initial quantity of 1-2 ng to a final quantity greater than 5 μ g. Amplified array elements are then purified using SEPHACRYL-400 (Amersham Pharmacia Biotech).

Purified array elements are immobilized on polymer-coated glass slides. Glass microscope slides (Corning) are cleaned by ultrasound in 0.1% SDS and acetone, with extensive distilled water washes between and after treatments. Glass slides are etched in 4% hydrofluoric acid (VWR Scientific Products Corporation (VWR), West Chester PA), washed extensively in distilled water, and coated with 0.05% aminopropyl silane (Sigma) in 95% ethanol. Coated slides are cured in a 110°C oven.

Array elements are applied to the coated glass substrate using a procedure described in US

Patent No. 5,807,522, incorporated herein by reference. 1 μ l of the array element DNA, at an average concentration of 100 ng/ μ l, is loaded into the open capillary printing element by a high-speed robotic apparatus. The apparatus then deposits about 5 nl of array element sample per slide.

Microarrays are UV-crosslinked using a STRATALINKER UV-crosslinker (Stratagene).

- 5 Microarrays are washed at room temperature once in 0.2% SDS and three times in distilled water. Non-specific binding sites are blocked by incubation of microarrays in 0.2% casein in phosphate buffered saline (PBS) (Tropix, Inc., Bedford MA) for 30 minutes at 60 °C followed by washes in 0.2% SDS and distilled water as before.

Hybridization

- 10 Hybridization reactions contain 9 μ l of sample mixture consisting of 0.2 μ g each of Cy3 and Cy5 labeled cDNA synthesis products in 5X SSC, 0.2% SDS hybridization buffer. The sample mixture is heated to 65 °C for 5 minutes and is aliquoted onto the microarray surface and covered with an 1.8 cm² coverslip. The arrays are transferred to a waterproof chamber having a cavity just slightly larger than a microscope slide. The chamber is kept at 100% humidity internally by the
- 15 addition of 140 μ l of 5X SSC in a corner of the chamber. The chamber containing the arrays is incubated for about 6.5 hours at 60 °C. The arrays are washed for 10 min at 45 °C in a first wash buffer (1X SSC, 0.1% SDS), three times for 10 minutes each at 45 °C in a second wash buffer (0.1X SSC), and dried.

Detection

- 20 Reporter-labeled hybridization complexes are detected with a microscope equipped with an Innova 70 mixed gas 10 W laser (Coherent, Inc., Santa Clara CA) capable of generating spectral lines at 488 nm for excitation of Cy3 and at 632 nm for excitation of Cy5. The excitation laser light is focused on the array using a 20X microscope objective (Nikon, Inc., Melville NY). The slide containing the array is placed on a computer-controlled X-Y stage on the microscope and raster-
- 25 scanned past the objective. The 1.8 cm x 1.8 cm array used in the present example is scanned with a resolution of 20 micrometers.

- In two separate scans, a mixed gas multiline laser excites the two fluorophores sequentially. Emitted light is split, based on wavelength, into two photomultiplier tube detectors (PMT R1477, Hamamatsu Photonics Systems, Bridgewater NJ) corresponding to the two fluorophores. Appropriate
- 30 filters positioned between the array and the photomultiplier tubes are used to filter the signals. The emission maxima of the fluorophores used are 565 nm for Cy3 and 650 nm for Cy5. Each array is typically scanned twice, one scan per fluorophore using the appropriate filters at the laser source, although the apparatus is capable of recording the spectra from both fluorophores simultaneously.

- The sensitivity of the scans is typically calibrated using the signal intensity generated by a
- 35 cDNA control species added to the sample mixture at a known concentration. A specific location on the array contains a complementary DNA sequence, allowing the intensity of the signal at that

location to be correlated with a weight ratio of hybridizing species of 1:100,000. When two samples from different sources (e.g., representing test and control cells), each labeled with a different fluorophore, are hybridized to a single array for the purpose of identifying genes that are differentially expressed, the calibration is done by labeling samples of the calibrating cDNA with the two fluorophores and adding identical amounts of each to the hybridization mixture.

The output of the photomultiplier tube is digitized using a 12-bit RTI-835H analog-to-digital (A/D) conversion board (Analog Devices, Inc., Norwood MA) installed in an IBM-compatible PC computer. The digitized data are displayed as an image where the signal intensity is mapped using a linear 20-color transformation to a pseudocolor scale ranging from blue (low signal) to red (high signal). The data is also analyzed quantitatively. Where two different fluorophores are excited and measured simultaneously, the data are first corrected for optical crosstalk (due to overlapping emission spectra) between the fluorophores using each fluorophore's emission spectrum.

A grid is superimposed over the fluorescence signal image such that the signal from each spot is centered in each element of the grid. The fluorescence signal within each element is then integrated to obtain a numerical value corresponding to the average intensity of the signal. The software used for signal analysis is the GEMTOOLS gene expression analysis program (Incyte).

VIII. Complementary Polynucleotides

Sequences complementary to the SYNT-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring SYNT. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of SYNT. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the SYNT-encoding transcript.

IX. Expression of SYNT

Expression and purification of SYNT is achieved using bacterial or virus-based expression systems. For expression of SYNT in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac (tac)* hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express SYNT upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of SYNT in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus

(AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding SYNT by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, SYNT is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from SYNT at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16). Purified SYNT obtained by these methods can be used directly in the assays shown in Examples X and XIV.

X. Demonstration of SYNT Activity

An SYNT activity assay measures aminoacylation of tRNA in the presence of a radiolabeled substrate. A cell-free extract depleted of endogenous aminoacyl-tRNA synthetase is prepared from Escherichia coli. SYNT, either biochemically purified or recombinantly produced, is added to the cell free extract. The cell-free extract is incubated with [¹⁴C]-labeled amino acid under conditions favorable for translation. Incorporation of the [¹⁴C]-labeled amino acid into acid-precipitable aminoacyl-tRNA is measured using a radioisotope counter. The amount of the [¹⁴C]-labeled amino acid incorporated into aminoacyl tRNA is proportional to the amount of SYNT activity. (See, for example, Ibba, M. et al. (1997) Science 278:1119-1122).

Alternatively, SYNT activity may be assayed as follows. SYNT, or biologically active fragments thereof, are labeled with ¹²⁵I Bolton-Hunter reagent. (See, e.g., Bolton et al. (1973) Biochem. J. 133:529.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled SYNT, washed, and any wells with labeled SYNT complex are assayed. Data obtained using different concentrations of SYNT are used to calculate values for the number, affinity, and association of SYNT with the candidate molecules.

XI. Functional Assays

SYNT function is assessed by expressing the sequences encoding SYNT at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include pCMV SPORT plasmid (Life Technologies) and pCR3.1 plasmid (Invitrogen), both of which contain the cytomegalovirus promoter. 5-10 μ g of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2 μ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of SYNT on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding SYNT and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding SYNT and other genes of interest can be analyzed by northern analysis or microarray techniques.

XII. Production of SYNT Specific Antibodies

SYNT substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the SYNT amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for

selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, supra, ch. 11.)

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (PE Biosystems) using Fmoc chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, supra.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for anti-peptide and anti-SYNT activity by, for example, binding the peptide or SYNT to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

10 XIII. Purification of Naturally Occurring SYNT Using Specific Antibodies

Naturally occurring or recombinant SYNT is substantially purified by immunoaffinity chromatography using antibodies specific for SYNT. An immunoaffinity column is constructed by covalently coupling anti-SYNT antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing SYNT are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of SYNT (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/SYNT binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and SYNT is collected.

20 XIV. Identification of Molecules Which Interact with SYNT

SYNT, or biologically active fragments thereof, are labeled with ¹²⁵I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) Biochem. J. 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled SYNT, washed, and any wells with labeled SYNT complex are assayed. Data obtained using different concentrations of SYNT are used to calculate values for the number, affinity, and association of SYNT with the candidate molecules.

Alternatively, molecules interacting with SYNT are analyzed using the yeast two-hybrid system as described in Fields, S. and O. Song (1989, Nature 340:245-246), or using commercially available kits based on the two-hybrid system, such as the MATCHMAKER system (Clontech).

SYNT may also be used in the PATHCALLING process (CuraGen Corp., New Haven CT) which employs the yeast two-hybrid system in a high-throughput manner to determine all interactions between the proteins encoded by two large libraries of genes (Nandabalan, K. et al. (2000) U.S. Patent No. 6,057,101).

35

Various modifications and variations of the described methods and systems of the invention

will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention
5 which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
1	16	1806212	SINTNOT13	268731R1 (HNT2NOT01), 410547R1 (BRSTNOT01), 642383F1 (BRSTNOT03), 642383R6 (BRSTNOT03), 1484778F6 (CORPNOT02), 1577276X13 (LNODNOT03), 1577276X14 (LNODNOT03), 1577276X16 (LNODNOT03), 1733808F6 (BRSTNOT08), 2057185R6 (BEPINOT01)
2	17	2083883	UTRSNOT08	1308703F1 (COLNFET02), 1337653F1 (COLNNOT13), 1430326F1 (SINTBST01), 1525049F6 (UCMCL5T01), 1525049T7 (UCMCL5T01), 1807032F6 (SINTNOT13), 2083883X31C1 (UTRSNOT08), 20838831X54C1, SAEA01919R1, SAEA00070F1
3	18	2454288	ENDANOT01	1334395F6 (COLNNOT13), 1607276F6 (LUNGNOT15), 1930074F6 (COLNTUT03), 1978977R6 (LUNGNOT03), 2054860R6 (BEPINOT01), 2454288H1 (ENDANOT01), 2791177H2 (COLNTUT16), 4049241H1 (SINTNOT18), 4787755H1 (BRATNOT03)
4	19	1513539	PANCTUT01	285705F1 (EOSIHET02), 515506F1 (MMLRIDT01), 515506R1 (MMLRIDT01), 723201R1 (SYNOOAT01), 726178R1 (SYNOOAT01), 906091R1 (COLNNOT08), 1492309T1 (PROSNON01), 1513539H1 (PANCTUT01), 2632033H1 (COLNTUT15)
5	20	2148623	BRAINOT09	655968H1 (EOSINOT03), 820407T1 (KERANOT02), 1383948F1 (BRAITUT08), 1383996T6 (BRAITUT08), 1441048R1 (THYRNOT03), 1579330F6 (DUODNOT01), 2148623H1 (BRAINOT09), 3170589H1 (BRSTNOT18), 3251591H1 (SEMVNOT03), 3618285H1 (EPIPNOT01), 3637486H1 (LUNGNOT30), g1524807
6	21	2579405	KIDNTUT13	2457R1 (U937NOT01), 270182R1 (HNT2NOT01), 730115H1 (LUNGNOT03), 991450H1 (COLNNOT11), 994055T1 (COLNNOT11), 1308496H1 (COLNFET02), 1808269F6 (SINTNOT13), 1875536T6 (LEUKNOT02), 1978055H1 (LUNGNOT03), 2579405H1 (KIDNTUT13), 2583944H1 (BRAITUT22)
7	22	2662427	ADRENOT08	283147F1 (CARDNOT01), 904594R2 (COLNNOT07), 2662427H1 (ADRENOT08), 2860066H1 (SININOT03), 3078653H1 (BRAIUNT01), 3852430F6 (BRAITUT12), 3852430T6 (BRAITUT12), SCDA09024V1, SCDA05143V1

Table 1 (cont.)

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
8	23	2844928	DRGLNOT01	1466601F1 (PANCUTUT02), 1662986F6 (BRSTNOT09), 1732880F6 (BRSTTUT08), 1804107F6 (SINTNOT13), 2521851F6 (BRAITUT21), 2662837X303B1 (ADRENOT08), 2844928H1 (DRGLNOT01), 2967532H1 (SCORNOT04), 3774082H1 (BRSTNOT25), 5871970H1 (COLTDIT04)
9	24	3231586	COTRNOT01	319277H1 (EOSIHET02), 920940H1 (RATRNOT02), 1445995H1 (PLACNOT02), 1517438F1 (PANCUTUT01), 1963042H1 (BRSTNOT04), 2023159H1 (CONNNOT01), 2932203H1 (UTRSNOR01), 3231586H1 (COTRNOT01), 3346128H1 (BRAITUT24), 91474393
10	25	3580770	293TF3T01	269986R1 (HNT2NOT01), 269986R6 (HNT2NOT01), 3580770H1 (293TF3T01), 3602575F6 (DRGTNOT01), 3602575T6 (DRGTNOT01), 91839725
11	26	3778612	BRSTNOT27	3778612F6 (BRSTNOT27), 3778612H1 (BRSTNOT27), 4063060T6 (BRAINOT21)
12	27	4574912	PROSTMT02	224714F1 (PANCNOT01), 452757F1 (TLYMNOT02), 1210705R7 (BRSTNOT02), 1437821F6 (PANCNOT08), 2700694F6 (OVARUTUT10), 3963634H1 (PROSNOT14), 3964731H1 (PROSNOT14), 4574912H1 (PROSTMT02), 5035739H1 (LIVRTUT13)
13	28	5630806	PLACFER01	1649584X20C1 (PROSTUT09), 1649855X12C1 (PROSTUT09), 1650519X14C1 (PROSTUT09), 2054061R6 (BEPINOT01), 2347805X304V1 (COLSUCT01), 3384595H1 (ESOGNOT04), 5630806H1 (PLACFER01)
14	29	5854855	FIBAUNT02	746434R1 (BRAITUT01), 815240R1 (OVARUTUT01), 2515988F6 (LIVRTUT04), 3105953F6 (BRSTTUT15), 5854855H1 (FIBAUNT02)
15	30	5993973	FTUBTUT02	270363H1 (HNT2NOT01), 1863382F6 (PROSNOT19), 2696339F6 (UTRSNOT12), 3500696H1 (PROSTUT13), 4247086H1 (BRABDIT01), 5195020H1 (LUNLTUT04), 5591818H1 (COLCDIT03), 6023466H1 (TESTNOT11)

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
1	1176	T7 T238 T543 S35 S68 S167 S175 T224 S241 T379 S396 T444 S453 S518 T552 S593 S673 T749 T769 T791 S978 S1042 T1083 T26 T41 S128 S153 T293 T327 S368 T557 S697 S714 S783 T925 S1082 T1136 S1153 Y111 Y666	N33 N333 N439 N882	Leucine-tRNA ligase domain T303-I870 Amino-transfer RNA synthetase N36-F88 N713-N723	similar to leucyl-tRNA synthetase [Arabidopsis thaliana] g2160156	MOTIFS BLAST-GenBank BLAST-DOMO PROFILES-SCAN BLAST-PRODROM BLIMPS-BLOCKS
2	739	S28 T132 S185 S282 S288 S307 T320 T352 S451 S661 T728 S732 S116 S163 S168 S261 T398 S436 T590 S633 T714 Y173 Y349 Y725 Y734	N102 N588 N619	AMP-binding domain S163-V629, N294-V345 E309-I320, T321-I329 Long-chain acyl-CoA synthetase P158-E280	fatty acid coenzyme A ligase 5 [Homo sapiens] g6174680	MOTIFS BLAST-GenBank BLAST-DOMO PROFILES-SCAN BLAST-PRODROM BLIMPS-BLOCKS HMMER-PFAM BLIMPS-PRINTS
3	589	S179 T18 T20 S40 S138 T193 T256 T298 T354 S419 S485 T570 S579 S5 T114 T174	N187	Phenylalanyl-tRNA synthetase P2-Q556, N293-I589 Synthetase b chain D59-L84, R355-Y374 I439-N465	phenylalanyl tRNA synthetase beta subunit [M. musculus] g4633656	MOTIFS BLAST-GenBank BLAST-PRODROM BLIMPS-PRODROM

Table 2 (cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
4	157	S26 S69 T131 T132 S108	N18	Signal peptide: M1-V66 Acetyl-CoA carboxylase carboxyl transferase alpha subunit signature: G128-I139	acetyl- coenzyme A synthetase [Penicillium chrysogenum] g265229	BLAST-GenBank SPSCAN BLOCKS-PRINTS MOTIFS
5	643	T308 T337 S402 S433 T454 S34 S101 T408 S530 S621	N9 N33 N90 N104 N259 N306 N431 N571 N634	Class II glutamine amidotransferase motif: L27-K36, G545-F560 (p<0.01) Transferase/ligase motif: R30-A224, S258-F356, P404-Q484	putative asparagine synthase [S. pombe] g3560144	BLAST-GenBank BLIMPS-BLOCKS BLAST-PRODOM MOTIFS
6	660	T282 S315 S329 T487 T596 S9 S276 T459 T624 Y230	N140	Class-I aminoacyl-tRNA synthetase signature: I74-M660, V183-L233, P201-L212, S519-N529 Arg-tRNA synthetase: K193-L224, V232-F245, T374-R395, H237-M660, S78-M660	arginyl-tRNA synthetase, ArgRS [Homo sapiens] g1217668	BLAST-GenBank HMMER-PFAM BLIMPS-BLOCKS PROFILESCAN BLIMPS-PFAM BLAST-PRODOM BLAST-DOMO MOTIFS

Table 2 (cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
7	725	S723 S80 S220 S238 T351 S371 T414 T465 S591 S641 T38 T45 T67 T165 S230 T252 T407 S413 S454 S499 S541 T554 T685 S688 T693 S719	N48 N472 N546 N552	Biotin-requiring enzyme signature: P652-F714, I657-R707 Biotin attachment site: K681 Acetyl-CoA biotin carboxyl carrier protein signature: F661-M675, V676-P689 Carbamoyl-PO4 synthase signature: K51-P490, E193-K494, F333-L340, L245-P290, Y270-K494, F331-G365	3-methyl- crotonyl CoA carboxylase homolog [C. elegans] g3876562	BLAST-GenBank HMMER-PFAM BLIMPS-BLOCKS PROFILES-SCAN BLIMPS-PRINTS BLAST-PRODROM BLAST-DOMO MOTIFS
8	644	S30 S253 S370 T457 T496 T593 T594 S43 T65 S207 S237 S269 S626 Y247 Y521	N282	AMP-binding domain signature: E79-V538, T101-V540, S240-L293, H255-L275, F262-H277, I260-K271, V570-R635 Ligase/synthetase signature: T356-R640	predicted long chain fatty acid CoA ligase [C. elegans] g4262587	BLAST-GenBank HMMER-PFAM BLIMPS-BLOCKS PROFILES-SCAN BLIMPS-PRINTS BLAST-PRODROM BLAST-DOMO MOTIFS
9	504	S122 S134 S428 S111 T349	N9 N39 N209 N426	Signal peptide: M1-A67 Aminotransferase class- II pyridoxal-PO4 attachment site: A104-R113		SPSCAN BLIMPS-BLOCKS MOTIFS

Table 2 (cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
10	489	S51 T150 T220 S465 S477 T90 S200 T250 T350 S433 S467 Y83		RNA methyl transferase SpuU family: P344-G354 CTP synthetase domain: K2-N455, M1-Y457, M335-Y457	CTP synthetase homolog (CTPsH) [Mus musculus] g1654185	BLAST-GenBank BLIMPS-PFAM BLAST-PRODOM BLAST-DOMO MOTIFS
11	258	S105 S115 S139		Signal peptide: M1-G43 ATP synthase signature: Y98-L129 S-adenosyl methionine synthetase signature: A209-V257		SPSCAN HMMER-PFAM PROFILES SCAN MOTIFS
12	555	S22 T72 T110 S132 S165 S190 S258 S342 S363 S446 S539 S31 T179 S212 S312 T318 S339 S380 S392 T393 T407 S428 T519 Y24		Aminoacyl tRNA synthetase signature: V73-A82, K305-N315 (p<1.1e-3) CysteinyI-tRNA synthetase signature: R66-A77, I103-V112, E230-C248, D261-E282, Y67-E282, H76-A494	putative cysteinyI- tRNA synthetase [S. pombe] g1044932	BLAST-GenBank BLIMPS-BLOCKS BLIMPS-PRINTS BLAST-PRODOM BLAST-DOMO MOTIFS
13	463	S54 S79 S124 S235 T253 T295 T411 T15 S366 Y84	N52	SuccinyI-CoA ligase family: I97-F111, V181-E193, L314-A343, L53-D367, D290-K442, L53-I463	ATP-specific succinyI-CoA synthetase beta subunit [Homo sapiens] g3766196	BLAST-GenBank BLIMPS-BLOCKS BLAST-PRODOM BLAST-DOMO MOTIFS

Table 2 (cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosyla- tion Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
14	399	T151 S179 S19 S72 T95 T151 S157 S191 S192 T217 T218 S260 T318 S337 S353		Signal peptide: M1-G42 Class-I aminoacyl-tRNA synthetase signature: P54-L64		SPSCAN MOTIFS
15	339	S187 T232 T266 S9 S31 T221 T231 S274 S276 T278 S324 S31 S36 T78 S223 S317 Y265		Phenylalanyl-tRNA synthetase family: V81-E261	similar to phenylalanyl- tRNA synthetase [C. elegans] g3876233	BLAST-GenBank BLAST-PROD MOTIFS

Table 3

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
16	153-197 600-644 1198-2042	Nervous (0.240) Reproductive (0.231) Hematopoietic/Immune (0.124)	Cancer (0.488) Inflammation/Trauma (0.306) Cell proliferation (0.198)	pINCY
17	163-207 1090-1134	Gastrointestinal (0.333) Reproductive (0.205) Hematopoietic/Immune (0.141)	Cancer (0.462) Inflammation/Trauma (0.372) Cell proliferation (0.103)	pINCY
18	601-645	Reproductive (0.216) Gastrointestinal (0.205) Nervous (0.148)	Cancer (0.477) Inflammation/Trauma (0.330) Cell proliferation (0.227)	PBLUESCRIPT
19	1-531 1634-2122	Reproductive (0.224) Nervous (0.165) Gastrointestinal (0.153)	Cancer (0.365) Inflammation (0.282) Cell proliferation (0.176)	pINCY
20	345-418 1272-1762 2039-2357	Cardiovascular (0.194) Reproductive (0.181) Nervous (0.153)	Cancer (0.514) Cell proliferation (0.194) Inflammation (0.181)	pINCY
21	1-109	Gastrointestinal (0.175) Nervous (0.175) Reproductive (0.175)	Cancer (0.427) Inflammation (0.252) Cell proliferation (0.243)	pINCY
22	2045-2317	Nervous (0.232) Reproductive (0.214) Cardiovascular (0.179)	Cancer (0.411) Inflammation (0.268) Trauma (0.161)	pINCY
23	1-508, 840-1183 1656-1741 1968-2254	Reproductive (0.266) Nervous (0.188) Endocrine (0.125)	Cancer (0.578) Inflammation (0.234) Cell proliferation (0.125)	pINCY
24		Reproductive (0.315) Gastrointestinal (0.150) Hematopoietic/Immune (0.150)	Cancer (0.433) Inflammation (0.283) Cell proliferation (0.126)	pINCY
25	1-151	Reproductive (0.333) Nervous (0.250) Hematopoietic (0.167)	Cancer (0.583) Inflammation (0.333) Cell proliferation (0.167)	pINCY

Table 3 (cont.)

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
26	1-214 567-970	Nervous (0.286) Reproductive (0.286) Cardiovascular (0.143) Endocrine (0.143)	Cancer (0.524) Inflammation (0.190) Trauma (0.190)	pINCY
27	1-206 1129-1206	Reproductive (0.243) Hematopoietic/Immune (0.143) Cardiovascular (0.129) Gastrointestinal (0.129) Nervous (0.129)	Cancer (0.457) Cell proliferation (0.229) Inflammation (0.186)	pINCY
28		Nervous (0.238) Gastrointestinal (0.190) Reproductive (0.190)	Cancer (0.429) Inflammation (0.274) Cell proliferation (0.131)	pINCY
29	1-234 1229-1477	Reproductive (0.331) Nervous (0.204) Gastrointestinal (0.116)	Cancer (0.508) Cell proliferation (0.188) Inflammation (0.182)	pINCY
30	1-871 1026-1138 1356-1660	Reproductive (0.378) Nervous (0.162) Gastrointestinal (0.135)	Cancer (0.649) Cell proliferation (0.189) Inflammation (0.189)	pINCY

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
16	SINTNOT13	Library was constructed using RNA isolated from ileum tissue obtained from a 25-year-old Asian female during a partial colectomy and temporary ileostomy. Pathology indicated moderately active chronic ulcerative colitis, involving colonic mucosa from the distal margin to the ascending colon. Family history included hyperlipidemia, malignant cervical neoplasm, viral hepatitis A, and depressive disorder.
17	UTRSNOT08	Library was constructed using RNA isolated from uterine tissue removed from a 35-year-old Caucasian female during a vaginal hysterectomy with dilation and curettage. Pathology indicated that the endometrium was secretory phase with a benign endometrial polyp 1 cm in diameter. The cervix showed mild chronic cervicitis. Family history included atherosclerotic coronary artery disease and type II diabetes.
18	ENDANOT01	Library was constructed using RNA isolated from aortic endothelial cell tissue from an explanted heart removed from a male during a heart transplant.
19	PANCTUT01	This library was constructed using RNA isolated from pancreatic tumor tissue removed from a 65-year-old Caucasian female. Pathology indicated invasive adenocarcinoma. Patient history included type II diabetes, osteoarthritis, cardiovascular disease, benign neoplasm in the large bowel, and a cataract. Family history included cardiovascular disease, type II diabetes, and stomach cancer.
20	BRAINOT09	This library was constructed using RNA isolated from brain tissue removed from a Caucasian male fetus, who died at 23 weeks' gestation.
21	KIDNTUT13	This library was constructed using RNA isolated from kidney tumor tissue removed from a 51-year-old Caucasian female. Pathology indicated renal cell carcinoma. Patient history included depressive disorder, hypoglycemia, and uterine endometriosis. Family history included calculus of the kidney, colon cancer, and type II diabetes.
22	ADRENOT08	This library was constructed using RNA isolated from adrenal tissue removed from a 20-year-old Caucasian male, who died from head trauma.

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
23	DRGLNOT01	This library was constructed using RNA isolated from dorsal root ganglion tissue removed from the low thoracic/high lumbar region of a 32-year-old Caucasian male who died from acute pulmonary edema and bronchopneumonia, bilateral pleural and pericardial effusions, and malignant lymphoma (natural killer cell type). Patient history included probable cytomegalovirus infection, hepatic congestion and steatosis, splenomegaly, hemorrhagic cystitis, thyroid hemorrhage, and Bell's palsy. Treatment included radiation therapy.
24	COTRNOT01	This library was constructed using RNA isolated from diseased transverse colon tissue obtained from a 26-year-old Caucasian male. Pathology indicated minimally active pancolitis with areas of focal severe colitis with perforation, consistent with Crohn's disease.
25	293TF3T01	This library was constructed using RNA isolated from a serum-starved transformed embryonal cell line (293-EBNA) derived from kidney epithelial tissue. The cells were transformed with adenovirus 5 DNA.
26	BRSTNOT27	This library was constructed using RNA isolated from breast tissue removed from a 57-year-old Caucasian female. Pathology for the associated tumor tissue indicated residual micro-spic infiltrating adenocarcinoma and extensive intraductal carcinoma. Immunoperoxidase stains for estrogen and progesterone receptors were positive. Patient history included benign hypertension, hyperlipidemia, cardiac dysrhythmia, a benign colon neoplasm, a solitary breast cyst, and a breast neoplasm. Family history included benign hypertension, acute leukemia, primary liver cancer, and lung cancer.
27	PROSTMT02	This library was constructed using RNA isolated from diseased prostate tissue removed from a 66-year-old Caucasian male. Pathology indicated adenofibromatous hyperplasia. Pathology from the associated tumor indicated adenocarcinoma. The patient presented with elevated prostate specific antigen (PSA) and induration. Family history included acute myocardial infarction, atherosclerotic coronary artery disease, type II diabetes, hyperlipidemia, and Jakob-Creutzfeldt disease.

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
28	PLACFER01	This library was constructed from RNA isolated from placental tissue from a Caucasian fetus, who died at 16 weeks' gestation from fetal demise and hydrocephalus. Patient history included umbilical cord wrapped around the head and shoulders. Serology was positive for anti-CMV. Family history included multiple pregnancies and live births, and an abortion.
29	FIBAUNT02	This library was constructed using RNA isolated from untreated aortic adventitial fibroblasts removed from a 65-year-old Caucasian female.
30	FTUBTUT02	This library was constructed using RNA isolated from fallopian tube tumor tissue removed from an 85-year-old Caucasian female. Pathology indicated poorly differentiated mixed endometrioid and serous adenocarcinoma. Pathology for the associated uterus tumor indicated focal and metastatic endometrioid adenocarcinoma and moderately differentiated invasive adenocarcinoma. Patient history included medullary carcinoma of the thyroid and myocardial infarction.

Table 5

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	PE Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	PE Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	PE Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25:3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. USA 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183:63-98; and Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value=1.06E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater fastx E value=1.0E-8 or less Full Length sequences: fastx score=100 or greater
BLIMPS	A BLocks IMProved Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff (1991) Nucleic Acids Res. 19:6565-6572; Henikoff, J.G. and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37:417-424.	Score=1000 or greater; Ratio of Score/Strength = 0.75 or larger; and, if applicable, Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) J. Mol. Biol. 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322.	Score=10-50 bits for PFAM hits, depending on individual protein families

Table 5 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, M. et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221.	Normalized quality score \geq GCG-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M.S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies.	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12:431-439.	Score=3.5 or greater
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

- 5 a) an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14,
- b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14,
- 10 c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, and
- 15 d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14.

20

2. An isolated polypeptide of claim 1 selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14.

25

3. An isolated polynucleotide encoding a polypeptide of claim 1.

4. An isolated polynucleotide encoding a polypeptide of claim 2.

5. An isolated polynucleotide of claim 4 selected from the group consisting of SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29.

30

6. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.

35

7. A cell transformed with a recombinant polynucleotide of claim 6.
8. A transgenic organism comprising a recombinant polynucleotide of claim 6.
- 5 9. A method for producing a polypeptide of claim 1, the method comprising:
- a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and
- 10 b) recovering the polypeptide so expressed.
10. An isolated antibody which specifically binds to a polypeptide of claim 1.
11. An isolated polynucleotide comprising a polynucleotide sequence selected from the
- 15 group consisting of:
- a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29,
- b) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a
- 20 polynucleotide sequence selected from the group consisting of SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29,
- c) a polynucleotide sequence complementary to a),
- d) a polynucleotide sequence complementary to b), and
- 25 e) an RNA equivalent of a)-d).
12. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 11.
- 30 13. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:
- a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization
- 35 complex is formed between said probe and said target polynucleotide or fragments thereof, and
- b) detecting the presence or absence of said hybridization complex, and, optionally, if

present, the amount thereof.

14. A method of claim 13, wherein the probe comprises at least 60 contiguous nucleotides.

5 15. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

 a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and

 b) detecting the presence or absence of said amplified target polynucleotide or fragment
10 thereof, and, optionally, if present, the amount thereof.

16. A composition comprising an effective amount of a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

15 17. A composition of claim 16, wherein the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14.

20 18. A method for treating a disease or condition associated with decreased expression of functional SYNT, comprising administering to a patient in need of such treatment the composition of claim 16.

 19. A method for screening a compound for effectiveness as an agonist of a polypeptide of
25 claim 1, the method comprising:

 a) exposing a sample comprising a polypeptide of claim 1 to a compound, and

 b) detecting agonist activity in the sample.

30 20. A composition comprising an agonist compound identified by a method of claim 19 and a pharmaceutically acceptable excipient.

 21. A method for treating a disease or condition associated with decreased expression of functional SYNT, comprising administering to a patient in need of such treatment a pharmaceutical composition of claim 20.

35 22. A method for screening a compound for effectiveness as an antagonist of a polypeptide

of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting antagonist activity in the sample.

5 23. A composition comprising an antagonist compound identified by a method of claim 22 and a pharmaceutically acceptable excipient.

 24. A method for treating a disease or condition associated with overexpression of functional SYNT, comprising administering to a patient in need of such treatment a composition of claim 23.

10

 25. A method of screening for a compound that specifically binds to the polypeptide of claim 1, said method comprising the steps of:

- a) combining the polypeptide of claim 1 with at least one test compound under suitable conditions, and
- 15 b) detecting binding of the polypeptide of claim 1 to the test compound, thereby identifying a compound that specifically binds to the polypeptide of claim 1.

 26. A method of screening for a compound that modulates the activity of the polypeptide of claim 1, said method comprising:

- 20 a) combining the polypeptide of claim 1 with at least one test compound under conditions permissive for the activity of the polypeptide of claim 1,
- b) assessing the activity of the polypeptide of claim 1 in the presence of the test compound, and
- c) comparing the activity of the polypeptide of claim 1 in the presence of the test compound
- 25 with the activity of the polypeptide of claim 1 in the absence of the test compound, wherein a change in the activity of the polypeptide of claim 1 in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide of claim 1.

 27. A method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 5, the method comprising:

- 30 a) exposing a sample comprising the target polynucleotide to a compound, and
- b) detecting altered expression of the target polynucleotide.

35 28. A method for assessing toxicity of a test compound, said method comprising:

- a) treating a biological sample containing nucleic acids with the test compound;

b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide of claim 11 under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence of a polynucleotide of claim

5 11 or fragment thereof;

c) quantifying the amount of hybridization complex; and

d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test

10 compound.

SEQUENCE LISTING

<110> INCYTE GENOMICS, INC.
 TANG, Y. Tom
 HILLMAN, Jennifer L.
 BANDMAN, Olga
 YUE, Henry
 BAUGHN, Mariah R.
 LAL, Preeti
 LU, Dyung Aina M.
 SHAH, Purvi
 AZIMZAI, Yalda

<120> HUMAN SYNTHETASES

<130> PF-0721 PCT

<140> To Be Assigned

<141> Herewith

<150> 60/144,992; 60,168,858

<151> 1999-07-22; 1999-12-02

<160> 30

<170> PERL Program

<210> 1

<211> 1176

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1806212CD1

<400> 1

Met	Ala	Glu	Arg	Lys	Gly	Thr	Ala	Lys	Val	Asp	Phe	Leu	Lys	Lys
1				5					10					15
Ile	Glu	Lys	Glu	Ile	Gln	Gln	Lys	Trp	Asp	Thr	Glu	Arg	Val	Phe
				20					25					30
Glu	Val	Asn	Ala	Ser	Asn	Leu	Glu	Lys	Gln	Thr	Ser	Lys	Gly	Lys
				35					40					45
Tyr	Phe	Val	Thr	Phe	Pro	Tyr	Pro	Tyr	Met	Asn	Gly	Arg	Leu	His
				50					55					60
Leu	Gly	His	Thr	Phe	Ser	Leu	Ser	Lys	Cys	Glu	Phe	Ala	Val	Gly
				65					70					75
Tyr	Gln	Arg	Leu	Lys	Gly	Lys	Cys	Cys	Leu	Phe	Pro	Phe	Gly	Leu
				80					85					90
His	Cys	Thr	Gly	Met	Pro	Ile	Lys	Ala	Cys	Ala	Asp	Lys	Leu	Lys
				95					100					105
Arg	Glu	Ile	Glu	Leu	Tyr	Gly	Cys	Pro	Pro	Asp	Phe	Pro	Asp	Glu
				110					115					120
Glu	Glu	Glu	Glu	Glu	Glu	Thr	Ser	Val	Lys	Thr	Glu	Asp	Ile	Ile
				125					130					135
Ile	Lys	Asp	Lys	Ala	Lys	Gly	Lys	Lys	Ser	Lys	Ala	Ala	Ala	Lys
				140					145					150
Ala	Gly	Ser	Ser	Lys	Tyr	Gln	Trp	Gly	Ile	Met	Lys	Ser	Leu	Gly
				155					160					165
Leu	Ser	Asp	Glu	Glu	Ile	Val	Lys	Phe	Ser	Glu	Ala	Glu	His	Trp
				170					175					180
Leu	Asp	Tyr	Phe	Pro	Pro	Leu	Ala	Ile	Gln	Asp	Leu	Lys	Arg	Met
				185					190					195

Gly	Leu	Lys	Val	Asp	Trp	Arg	Arg	Ser	Phe	Ile	Thr	Thr	Asp	Val
				200					205					210
Asn	Pro	Tyr	Tyr	Asp	Ser	Phe	Val	Arg	Trp	Gln	Phe	Leu	Thr	Leu
				215					220					225
Arg	Glu	Arg	Asn	Lys	Ile	Lys	Phe	Gly	Lys	Arg	Tyr	Thr	Ile	Tyr
				230					235					240
Ser	Pro	Lys	Asp	Gly	Gln	Pro	Cys	Met	Asp	His	Asp	Arg	Gln	Thr
				245					250					255
Gly	Glu	Gly	Val	Gly	Pro	Gln	Glu	Tyr	Thr	Leu	Leu	Lys	Leu	Lys
				260					265					270
Val	Leu	Glu	Pro	Tyr	Pro	Ser	Lys	Leu	Ser	Gly	Leu	Lys	Gly	Lys
				275					280					285
Asn	Ile	Phe	Leu	Val	Ala	Ala	Thr	Leu	Arg	Pro	Glu	Thr	Met	Phe
				290					295					300
Gly	Gln	Thr	Asn	Cys	Trp	Val	Arg	Pro	Asp	Met	Lys	Tyr	Ile	Gly
				305					310					315
Phe	Glu	Thr	Val	Asn	Gly	Asp	Ile	Phe	Ile	Cys	Thr	Gln	Lys	Ala
				320					325					330
Ala	Arg	Asn	Met	Ser	Tyr	Gln	Gly	Phe	Thr	Lys	Asp	Asn	Gly	Val
				335					340					345
Val	Pro	Val	Val	Lys	Glu	Leu	Met	Gly	Glu	Glu	Ile	Leu	Gly	Ala
				350					355					360
Ser	Leu	Ser	Ala	Pro	Leu	Thr	Ser	Tyr	Lys	Val	Ile	Tyr	Val	Leu
				365					370					375
Pro	Met	Leu	Thr	Ile	Lys	Glu	Asp	Lys	Gly	Thr	Gly	Val	Val	Thr
				380					385					390
Ser	Val	Pro	Ser	Asp	Ser	Pro	Asp	Asp	Ile	Ala	Ala	Leu	Arg	Asp
				395					400					405
Leu	Lys	Lys	Lys	Gln	Ala	Leu	Arg	Ala	Lys	Tyr	Gly	Ile	Arg	Asp
				410					415					420
Asp	Met	Val	Leu	Pro	Phe	Glu	Pro	Val	Pro	Val	Ile	Glu	Ile	Pro
				425					430					435
Gly	Phe	Gly	Asn	Leu	Ser	Ala	Val	Thr	Ile	Cys	Asp	Glu	Leu	Lys
				440					445					450
Ile	Gln	Ser	Gln	Asn	Asp	Arg	Glu	Lys	Leu	Ala	Glu	Ala	Lys	Glu
				455					460					465
Lys	Ile	Tyr	Leu	Lys	Gly	Phe	Tyr	Glu	Gly	Ile	Met	Leu	Val	Asp
				470					475					480
Gly	Phe	Lys	Gly	Gln	Lys	Val	Gln	Asp	Val	Lys	Lys	Thr	Ile	Gln
				485					490					495
Lys	Lys	Met	Ile	Asp	Ala	Gly	Asp	Ala	Leu	Ile	Tyr	Met	Glu	Pro
				500					505					510
Glu	Lys	Gln	Val	Met	Ser	Arg	Ser	Ser	Asp	Glu	Cys	Val	Val	Ala
				515					520					525
Leu	Cys	Asp	Gln	Trp	Tyr	Leu	Asp	Tyr	Gly	Glu	Glu	Asn	Trp	Lys
				530					535					540
Lys	Gln	Thr	Ser	Gln	Cys	Leu	Lys	Asn	Leu	Glu	Thr	Phe	Cys	Glu
				545					550					555
Glu	Thr	Arg	Arg	Asn	Phe	Glu	Ala	Thr	Leu	Gly	Trp	Leu	Gln	Glu
				560					565					570
His	Ala	Cys	Ser	Arg	Thr	Tyr	Gly	Leu	Gly	Thr	His	Leu	Pro	Trp
				575					580					585
Asp	Glu	Gln	Trp	Leu	Ile	Glu	Ser	Leu	Ser	Asp	Ser	Thr	Ile	Tyr
				590					595					600
Met	Ala	Phe	Tyr	Thr	Val	Ala	His	Leu	Leu	Gln	Gly	Gly	Asn	Leu
				605					610					615
His	Gly	Gln	Ala	Glu	Ser	Pro	Leu	Gly	Ile	Arg	Pro	Gln	Gln	Met
				620					625					630
Thr	Lys	Glu	Val	Trp	Asp	Tyr	Val	Phe	Phe	Lys	Glu	Ala	Pro	Phe
				635					640					645
Pro	Lys	Thr	Gln	Ile	Ala	Lys	Glu	Lys	Leu	Asp	Gln	Leu	Lys	Gln
				650					655					660
Glu	Phe	Glu	Phe	Trp	Tyr	Pro	Val	Asp	Leu	Arg	Val	Ser	Gly	Lys
				665					670					675
Asp	Leu	Val	Pro	Asn	His	Leu	Ser	Tyr	Tyr	Leu	Tyr	Asn	His	Val

Ala Met Trp Pro	680	685	690
Glu Gln Ser Asp Lys Trp Pro Thr Ala Val Arg	695	700	705
Ala Asn Gly His Leu Leu Leu Asn Ser Glu Lys Met Ser Lys Ser	710	715	720
Thr Gly Asn Phe Leu Thr Leu Thr Gln Ala Ile Asp Lys Phe Ser	725	730	735
Ala Asp Gly Met Arg Leu Ala Leu Ala Asp Ala Gly Asp Thr Val	740	745	750
Glu Asp Ala Asn Phe Val Glu Ala Met Ala Asp Ala Gly Ile Leu	755	760	765
Arg Leu Tyr Thr Trp Val Glu Trp Val Lys Glu Met Val Ala Asn	770	775	780
Trp Asp Ser Leu Arg Ser Gly Pro Ala Ser Thr Phe Asn Asp Arg	785	790	795
Val Phe Ala Ser Glu Leu Asn Ala Gly Ile Ile Lys Thr Asp Gln	800	805	810
Asn Tyr Glu Lys Met Met Phe Lys Glu Ala Leu Lys Thr Gly Phe	815	820	825
Phe Glu Phe Gln Ala Ala Lys Asp Lys Tyr Arg Glu Leu Ala Val	830	835	840
Glu Gly Met His Arg Glu Leu Val Phe Arg Phe Ile Glu Val Gln	845	850	855
Thr Leu Leu Leu Ala Pro Phe Cys Pro His Leu Cys Glu His Ile	860	865	870
Trp Thr Leu Leu Gly Lys Pro Asp Ser Ile Met Asn Ala Ser Trp	875	880	885
Pro Val Ala Gly Pro Val Asn Glu Val Leu Ile His Ser Ser Gln	890	895	900
Tyr Leu Met Glu Val Thr His Asp Leu Arg Leu Arg Leu Lys Asn	905	910	915
Tyr Met Met Pro Ala Lys Gly Lys Lys Thr Asp Lys Gln Pro Leu	920	925	930
Gln Lys Pro Ser His Cys Thr Ile Tyr Val Ala Lys Asn Tyr Pro	935	940	945
Pro Trp Gln His Thr Thr Leu Ser Val Leu Arg Lys His Phe Glu	950	955	960
Ala Asn Asn Gly Lys Leu Pro Asp Asn Lys Val Ile Ala Ser Glu	965	970	975
Leu Gly Ser Met Pro Glu Leu Lys Lys Tyr Met Lys Lys Val Met	980	985	990
Pro Phe Val Ala Met Ile Lys Glu Asn Leu Glu Lys Met Gly Pro	995	1000	1005
Arg Ile Leu Asp Leu Gln Leu Glu Phe Asp Glu Lys Ala Val Leu	1010	1015	1020
Met Glu Asn Ile Val Tyr Leu Thr Asn Ser Leu Glu Leu Glu His	1025	1030	1035
Ile Glu Val Lys Phe Ala Ser Glu Ala Glu Asp Lys Ile Arg Glu	1040	1045	1050
Asp Cys Cys Pro Gly Lys Pro Leu Asn Val Phe Arg Ile Glu Pro	1055	1060	1065
Gly Val Ser Val Ser Leu Val Asn Pro Gln Pro Ser Asn Gly His	1070	1075	1080
Phe Ser Thr Lys Ile Glu Ile Arg Gln Gly Asp Asn Cys Asp Ser	1085	1090	1095
Ile Ile Arg Arg Leu Met Lys Met Asn Arg Gly Ile Lys Asp Leu	1100	1105	1110
Ser Lys Val Lys Leu Met Arg Phe Asp Asp Pro Leu Leu Gly Pro	1115	1120	1125
Arg Arg Val Pro Val Leu Gly Lys Glu Tyr Thr Glu Lys Thr Pro	1130	1135	1140
Ile Ser Glu His Ala Val Phe Asn Val Asp Leu Met Ser Lys Lys	1145	1150	1155
Ile His Leu Thr Glu Asn Gly Ile Arg Val Asp Ile Gly Asp Thr	1160	1165	1170

Ile Ile Tyr Leu Val His
1175

<210> 2

<211> 739

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2083883CD1

<400> 2

Met	Asp	Ala	Leu	Lys	Pro	Pro	Cys	Leu	Trp	Arg	Asn	His	Glu	Arg	1	5	10	15
Gly	Lys	Lys	Asp	Arg	Asp	Ser	Cys	Gly	Arg	Lys	Asn	Ser	Glu	Pro	20	25	30	35
Gly	Ser	Pro	His	Ser	Leu	Glu	Ala	Leu	Arg	Asp	Ala	Ala	Pro	Ser	40	45	50	55
Gln	Gly	Leu	Asn	Phe	Leu	Leu	Leu	Phe	Thr	Lys	Met	Leu	Phe	Ile	60	65	70	75
Phe	Asn	Phe	Leu	Phe	Ser	Pro	Leu	Pro	Thr	Pro	Ala	Leu	Ile	Cys	80	85	90	95
Ile	Leu	Thr	Phe	Gly	Ala	Ala	Ile	Phe	Leu	Trp	Leu	Ile	Thr	Arg	100	105	110	115
Pro	Gln	Pro	Val	Leu	Pro	Leu	Leu	Asp	Leu	Asn	Asn	Gln	Ser	Val	120	125	130	135
Gly	Ile	Glu	Gly	Gly	Ala	Arg	Lys	Gly	Val	Ser	Gln	Lys	Asn	Asn	140	145	150	155
Asp	Leu	Thr	Ser	Cys	Cys	Phe	Ser	Asp	Ala	Lys	Thr	Met	Tyr	Glu	160	165	170	175
Val	Phe	Gln	Arg	Gly	Leu	Ala	Val	Ser	Asp	Asn	Gly	Pro	Cys	Leu	180	185	190	195
Gly	Tyr	Arg	Lys	Pro	Asn	Gln	Pro	Tyr	Arg	Trp	Leu	Ser	Tyr	Lys	200	205	210	215
Gln	Val	Ser	Asp	Arg	Ala	Glu	Tyr	Leu	Gly	Ser	Cys	Leu	Leu	His	220	225	230	235
Lys	Gly	Tyr	Lys	Ser	Ser	Pro	Asp	Gln	Phe	Val	Gly	Ile	Phe	Ala	240	245	250	255
Gln	Asn	Arg	Pro	Glu	Trp	Ile	Ile	Ser	Glu	Leu	Ala	Cys	Tyr	Thr	260	265	270	275
Tyr	Ser	Met	Val	Ala	Val	Pro	Leu	Tyr	Asp	Thr	Leu	Gly	Pro	Glu	280	285	290	295
Ala	Ile	Val	His	Ile	Val	Asn	Lys	Ala	Asp	Ile	Ala	Val	Val	Ile	300	305	310	315
Cys	Asp	Thr	Pro	Gln	Lys	Ala	Leu	Val	Leu	Ile	Gly	Asn	Val	Glu	320	325	330	335
Lys	Gly	Phe	Thr	Pro	Ser	Leu	Lys	Val	Ile	Ile	Leu	Met	Asp	Pro	340	345	350	355
Phe	Asp	Asp	Asp	Leu	Lys	Gln	Arg	Gly	Glu	Lys	Ser	Gly	Ile	Glu	360	365	370	375
Ile	Leu	Ser	Leu	Tyr	Asp	Ala	Glu	Asn	Leu	Gly	Lys	Glu	His	Phe	380	385	390	395
Arg	Lys	Pro	Val	Pro	Pro	Ser	Pro	Glu	Asp	Leu	Ser	Val	Ile	Cys	400	405	410	415
Phe	Thr	Ser	Gly	Thr	Thr	Gly	Asp	Pro	Lys	Gly	Ala	Met	Ile	Thr	420	425	430	435
His	Gln	Asn	Ile	Val	Ser	Asn	Ala	Ala	Ala	Phe	Leu	Lys	Cys	Val	440	445	450	455
Glu	His	Ala	Tyr	Glu	Pro	Thr	Pro	Asp	Asp	Val	Ala	Ile	Ser	Tyr	460	465	470	475
Leu	Pro	Leu	Ala	His	Met	Phe	Glu	Arg	Ile	Val	Gln	Ala	Val	Val	480	485	490	495
Tyr	Ser	Cys	Gly	Ala	Arg	Val	Gly	Phe	Phe	Gln	Gly	Asp	Ile	Arg	500	505	510	515
Leu	Leu	Ala	Asp	Asp	Met	Lys	Thr	Leu	Lys	Pro	Thr	Leu	Phe	Pro	520	525	530	535

	395		400		405
Ala Val Pro Arg	Leu Leu Asn Arg Ile Tyr Asp Lys Val Gln Asn				
	410		415		420
Glu Ala Lys Thr	Pro Leu Lys Lys Phe Leu Leu Lys Leu Ala Val				
	425		430		435
Ser Ser Lys Phe	Lys Glu Leu Gln Lys Gly Ile Ile Arg His Asp				
	440		445		450
Ser Phe Trp Asp	Lys Leu Ile Phe Ala Lys Ile Gln Asp Ser Leu				
	455		460		465
Gly Gly Arg Val	Arg Val Ile Val Thr Gly Ala Ala Pro Met Ser				
	470		475		480
Thr Ser Val Met	Thr Phe Phe Arg Ala Ala Met Gly Cys Gln Val				
	485		490		495
Tyr Glu Ala Tyr	Gly Gln Thr Glu Cys Thr Gly Gly Cys Thr Phe				
	500		505		510
Thr Leu Pro Gly	Asp Trp Thr Ser Gly His Val Gly Val Pro Leu				
	515		520		525
Ala Cys Asn Tyr	Val Lys Leu Glu Asp Val Ala Asp Met Asn Tyr				
	530		535		540
Phe Thr Val Asn	Asn Glu Gly Glu Val Cys Ile Lys Gly Thr Asn				
	545		550		555
Val Phe Lys Gly	Tyr Leu Lys Asp Pro Glu Lys Thr Gln Glu Ala				
	560		565		570
Leu Asp Ser Asp	Gly Trp Leu His Thr Gly Asp Ile Gly Arg Trp				
	575		580		585
Leu Pro Asn Gly	Thr Leu Lys Ile Ile Asp Arg Lys Lys Asn Ile				
	590		595		600
Phe Lys Leu Ala	Gln Gly Glu Tyr Ile Ala Pro Glu Lys Ile Glu				
	605		610		615
Asn Ile Tyr Asn	Arg Ser Gln Pro Val Leu Gln Ile Phe Val His				
	620		625		630
Gly Glu Ser Leu	Arg Ser Ser Leu Val Gly Val Val Val Pro Asp				
	635		640		645
Thr Asp Val Leu	Pro Ser Phe Ala Ala Lys Leu Gly Val Lys Gly				
	650		655		660
Ser Phe Glu Glu	Leu Cys Gln Asn Gln Val Val Arg Glu Ala Ile				
	665		670		675
Leu Glu Asp Leu	Gln Lys Ile Gly Lys Glu Ser Gly Leu Lys Thr				
	680		685		690
Phe Glu Gln Val	Lys Ala Ile Phe Leu His Pro Glu Pro Phe Ser				
	695		700		705
Ile Glu Asn Gly	Leu Leu Thr Pro Thr Leu Lys Ala Lys Arg Gly				
	710		715		720
Glu Leu Ser Lys	Tyr Phe Arg Thr Gln Ile Asp Ser Leu Tyr Glu				
	725		730		735
His Ile Gln Asp					

<210> 3

<211> 589

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2454288CD1

<400> 3

Met Pro Thr Val	Ser Val Lys Arg Asp Leu Leu Phe Gln Ala Leu	
1	5	10
Gly Arg Thr Tyr	Thr Asp Glu Glu Phe Asp Glu Leu Cys Phe Glu	
	20	25
Phe Gly Leu Glu	Leu Asp Glu Ile Thr Ser Glu Lys Glu Ile Ile	
	35	40
Ser Lys Glu Gln	Gly Asn Val Lys Ala Ala Gly Ala Ser Asp Val	
	50	55
		60

Val	Leu	Tyr	Lys	Ile	Asp	Val	Pro	Ala	Asn	Arg	Tyr	Asp	Leu	Leu
				65					70					75
Cys	Leu	Glu	Gly	Leu	Val	Arg	Gly	Leu	Gln	Val	Phe	Lys	Glu	Arg
				80					85					90
Ile	Lys	Ala	Pro	Val	Tyr	Lys	Arg	Val	Met	Pro	Asp	Gly	Lys	Ile
				95					100					105
Gln	Lys	Leu	Ile	Ile	Thr	Glu	Glu	Thr	Ala	Lys	Ile	Arg	Pro	Phe
				110					115					120
Ala	Val	Ala	Ala	Val	Leu	Arg	Asn	Ile	Lys	Phe	Thr	Lys	Asp	Arg
				125					130					135
Tyr	Asp	Ser	Phe	Ile	Glu	Leu	Gln	Glu	Lys	Leu	His	Gln	Asn	Ile
				140					145					150
Cys	Arg	Lys	Arg	Ala	Leu	Val	Ala	Ile	Gly	Thr	His	Asp	Leu	Asp
				155					160					165
Thr	Leu	Ser	Gly	Pro	Phe	Thr	Tyr	Thr	Ala	Lys	Arg	Pro	Ser	Asp
				170					175					180
Ile	Lys	Phe	Lys	Pro	Leu	Asn	Lys	Thr	Lys	Glu	Tyr	Thr	Ala	Cys
				185					190					195
Glu	Leu	Met	Asn	Ile	Tyr	Lys	Thr	Asp	Asn	His	Leu	Lys	His	Tyr
				200					205					210
Leu	His	Ile	Ile	Glu	Asn	Lys	Pro	Leu	Tyr	Pro	Val	Ile	Tyr	Asp
				215					220					225
Ser	Asn	Gly	Val	Val	Leu	Ser	Met	Pro	Pro	Ile	Ile	Asn	Gly	Asp
				230					235					240
His	Ser	Arg	Ile	Thr	Val	Asn	Thr	Arg	Asn	Ile	Phe	Ile	Glu	Cys
				245					250					255
Thr	Gly	Thr	Asp	Phe	Thr	Lys	Ala	Lys	Ile	Val	Leu	Asp	Ile	Ile
				260					265					270
Val	Thr	Met	Phe	Ser	Glu	Tyr	Cys	Glu	Asn	Gln	Phe	Thr	Val	Glu
				275					280					285
Ala	Ala	Glu	Val	Val	Phe	Pro	Asn	Gly	Lys	Ser	His	Thr	Phe	Pro
				290					295					300
Glu	Leu	Ala	Tyr	Arg	Lys	Glu	Met	Val	Arg	Ala	Asp	Leu	Ile	Asn
				305					310					315
Lys	Lys	Val	Gly	Ile	Arg	Glu	Thr	Pro	Glu	Asn	Leu	Ala	Lys	Leu
				320					325					330
Leu	Thr	Arg	Met	Tyr	Leu	Lys	Ser	Glu	Val	Ile	Gly	Asp	Gly	Asn
				335					340					345
Gln	Ile	Glu	Ile	Glu	Ile	Pro	Pro	Thr	Arg	Ala	Asp	Ile	Ile	His
				350					355					360
Ala	Cys	Asp	Ile	Val	Glu	Asp	Ala	Ala	Ile	Ala	Tyr	Gly	Tyr	Asn
				365					370					375
Asn	Ile	Gln	Met	Thr	Leu	Pro	Lys	Thr	Tyr	Thr	Ile	Ala	Asn	Gln
				380					385					390
Phe	Pro	Leu	Asn	Lys	Leu	Thr	Glu	Leu	Leu	Arg	His	Asp	Met	Ala
				395					400					405
Ala	Ala	Gly	Phe	Thr	Glu	Ala	Leu	Thr	Phe	Ala	Leu	Cys	Ser	Gln
				410					415					420
Glu	Asp	Ile	Ala	Asp	Lys	Leu	Gly	Val	Asp	Ile	Ser	Ala	Thr	Lys
				425					430					435
Ala	Val	His	Ile	Ser	Asn	Pro	Lys	Thr	Ala	Glu	Phe	Gln	Val	Ala
				440					445					450
Arg	Thr	Thr	Leu	Leu	Pro	Gly	Leu	Leu	Lys	Thr	Ile	Ala	Ala	Asn
				455					460					465
Arg	Lys	Met	Pro	Leu	Pro	Leu	Lys	Leu	Phe	Glu	Ile	Ser	Asp	Ile
				470					475					480
Val	Ile	Lys	Asp	Ser	Asn	Thr	Asp	Val	Gly	Ala	Lys	Asn	Tyr	Arg
				485					490					495
His	Leu	Cys	Ala	Val	Tyr	Tyr	Asn	Lys	Asn	Pro	Gly	Phe	Glu	Ile
				500					505					510
Ile	His	Gly	Leu	Leu	Asp	Arg	Ile	Met	Gln	Leu	Leu	Asp	Val	Pro
				515					520					525
Pro	Gly	Glu	Asp	Lys	Gly	Gly	Tyr	Val	Ile	Lys	Ala	Ser	Glu	Gly
				530					535					540
Pro	Ala	Phe	Phe	Pro	Gly	Arg	Cys	Ala	Glu	Ile	Phe	Ala	Arg	Gly

Gln Ser Val Gly	545	Leu Gly Val Leu	550	His Pro Asp Val Ile Thr	555
Lys Phe Glu Leu Thr	560	Met Pro Cys Ser	565	Ser Leu Glu Ile Asn Ile	570
Gly Pro Phe Leu	575		580		585

<210> 4
 <211> 157
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1513539CD1

<400> 4

Met Lys Leu Lys Cys	Ile Phe Gly Phe Ala	Thr Lys Glu Thr Ser
1 5	10	15
Cys Tyr Asn Val Thr	Asn Ile Gly Phe Lys	Ser Pro Ser Asp Phe
20	25	30
Trp Gln Ser Val His	Ser Thr Leu Pro Arg	Glu Leu Ala Pro Cys
35	40	45
Leu Val Phe Asn Thr	Ser Pro Asn Leu Ala	Leu Phe Ser Ala Ala
50	55	60
Phe Ala Phe Ile Val	Val Lys Asp Ser Ala	Gly Asp Ser Asp Val
65	70	75
Val Val Gln Glu Leu	Lys Ser Met Val Ala	Thr Lys Ile Ala Lys
80	85	90
Tyr Ala Val Pro Asp	Glu Ile Leu Val Val	Lys Arg Leu Pro Lys
95	100	105
Thr Arg Ser Gly Lys	Val Met Arg Arg Leu	Leu Arg Lys Ile Ile
110	115	120
Thr Ser Glu Ala Gln	Glu Leu Gly Asp Thr	Thr Thr Leu Glu Asp
125	130	135
Pro Ser Ile Ile Ala	Glu Ile Leu Ser Val	Tyr Gln Lys Cys Lys
140	145	150
Asp Lys Gln Ala Ala	Lys	
155		

<210> 5
 <211> 643
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2148623CD1

<400> 5

Met Cys Gly Ile Cys	Cys Ser Val Asn Phe	Ser Ala Glu His Phe
1 5	10	15
Ser Gln Asp Leu Lys	Glu Asp Leu Leu Tyr	Asn Leu Lys Gln Arg
20	25	30
Gly Pro Asn Ser Ser	Lys Gln Leu Leu Lys	Ser Asp Val Asn Tyr
35	40	45
Gln Cys Leu Phe Ser	Ala His Val Leu His	Leu Arg Gly Val Leu
50	55	60
Thr Thr Gln Pro Val	Glu Asp Glu Arg Gly	Asn Val Phe Leu Trp
65	70	75
Asn Gly Glu Ile Phe	Ser Gly Ile Lys Val	Glu Ala Glu Glu Asn
80	85	90
Asp Thr Gln Ile Leu	Phe Asn Tyr Leu Ser	Ser Cys Lys Asn Glu
95	100	105
Ser Glu Ile Leu Ser	Leu Phe Ser Glu Val	Gln Gly Pro Trp Ser
110	115	120

Phe	Ile	Tyr	Tyr	Gln	Ala	Ser	Ser	His	Tyr	Leu	Trp	Phe	Gly	Arg
				125					130					135
Asp	Phe	Phe	Gly	Arg	Arg	Ser	Leu	Leu	Trp	His	Phe	Ser	Asn	Leu
				140					145					150
Gly	Lys	Ser	Phe	Cys	Leu	Ser	Ser	Val	Gly	Thr	Gln	Thr	Ser	Gly
				155					160					165
Leu	Ala	Asn	Gln	Trp	Gln	Glu	Val	Pro	Ala	Ser	Gly	Leu	Phe	Arg
				170					175					180
Ile	Asp	Leu	Lys	Ser	Thr	Val	Ile	Ser	Arg	Cys	Ile	Ile	Leu	Gln
				185					190					195
Leu	Tyr	Pro	Trp	Lys	Tyr	Ile	Ser	Arg	Glu	Asn	Ile	Ile	Glu	Glu
				200					205					210
Asn	Val	Asn	Ser	Leu	Ser	Gln	Ile	Ser	Ala	Asp	Leu	Pro	Ala	Phe
				215					220					225
Val	Ser	Val	Val	Ala	Asn	Glu	Ala	Lys	Leu	Tyr	Leu	Glu	Lys	Pro
				230					235					240
Val	Val	Pro	Leu	Asn	Met	Met	Leu	Pro	Gln	Ala	Ala	Leu	Glu	Thr
				245					250					255
His	Cys	Ser	Asn	Ile	Ser	Asn	Val	Pro	Pro	Thr	Arg	Glu	Ile	Leu
				260					265					270
Gln	Val	Phe	Leu	Thr	Asp	Val	His	Met	Lys	Glu	Val	Ile	Gln	Gln
				275					280					285
Phe	Ile	Asp	Val	Leu	Ser	Val	Ala	Val	Lys	Lys	Arg	Val	Leu	Cys
				290					295					300
Leu	Pro	Arg	Asp	Glu	Asn	Leu	Thr	Ala	Asn	Glu	Val	Leu	Lys	Thr
				305					310					315
Cys	Asp	Arg	Lys	Ala	Asn	Val	Ala	Ile	Leu	Phe	Ser	Gly	Gly	Ile
				320					325					330
Asp	Ser	Met	Val	Ile	Ala	Thr	Leu	Ala	Asp	Arg	His	Ile	Pro	Leu
				335					340					345
Asp	Glu	Pro	Ile	Asp	Leu	Leu	Asn	Val	Ala	Phe	Ile	Ala	Glu	Glu
				350					355					360
Lys	Thr	Met	Pro	Thr	Thr	Phe	Asn	Arg	Glu	Gly	Asn	Lys	Gln	Lys
				365					370					375
Asn	Lys	Cys	Glu	Ile	Pro	Ser	Glu	Glu	Phe	Ser	Lys	Asp	Val	Ala
				380					385					390
Ala	Ala	Ala	Ala	Asp	Ser	Pro	Asn	Lys	His	Val	Ser	Val	Pro	Asp
				395					400					405
Arg	Ile	Thr	Gly	Arg	Ala	Gly	Leu	Lys	Glu	Leu	Gln	Ala	Val	Ser
				410					415					420
Pro	Ser	Arg	Ile	Trp	Asn	Phe	Val	Glu	Ile	Asn	Val	Ser	Met	Glu
				425					430					435
Glu	Leu	Gln	Lys	Leu	Arg	Arg	Thr	Arg	Ile	Cys	His	Leu	Ile	Arg
				440					445					450
Pro	Leu	Asp	Thr	Val	Leu	Asp	Asp	Ser	Ile	Gly	Cys	Ala	Val	Trp
				455					460					465
Phe	Ala	Ser	Arg	Gly	Ile	Gly	Trp	Leu	Val	Ala	Gln	Glu	Gly	Val
				470					475					480
Lys	Ser	Tyr	Gln	Ser	Asn	Ala	Lys	Val	Val	Leu	Thr	Gly	Ile	Gly
				485					490					495
Ala	Asp	Glu	Gln	Leu	Ala	Gly	Tyr	Ser	Arg	His	Arg	Val	Arg	Phe
				500					505					510
Gln	Ser	His	Gly	Leu	Glu	Gly	Leu	Asn	Lys	Glu	Ile	Met	Met	Glu
				515					520					525
Leu	Gly	Arg	Ile	Ser	Ser	Arg	Asn	Leu	Gly	Arg	Asp	Asp	Arg	Val
				530					535					540
Ile	Gly	Asp	His	Gly	Lys	Glu	Ala	Arg	Phe	Pro	Phe	Leu	Asp	Glu
				545					550					555
Asn	Val	Val	Ser	Phe	Leu	Asn	Ser	Leu	Pro	Ile	Trp	Glu	Lys	Ala
				560					565					570
Asn	Leu	Thr	Leu	Pro	Arg	Gly	Ile	Gly	Glu	Lys	Leu	Leu	Leu	Arg
				575					580					585
Leu	Ala	Ala	Val	Glu	Leu	Gly	Leu	Thr	Ala	Ser	Ala	Leu	Leu	Pro
				590					595					600
Lys	Arg	Ala	Met	Gln	Phe	Gly	Ser	Arg	Ile	Ala	Lys	Met	Glu	Lys

	605		610		615
Ile Asn Glu Lys	Ala Ser Asp Lys Cys	Gly Arg Leu Gln Ile	Met		
	620		625		630
Ser Leu Glu Asn	Leu Ser Ile Glu Lys	Glu Thr Lys Leu			
	635		640		

<210> 6
 <211> 660
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2579405CD1

<400> 6
 Met Asp Ile Leu Val Ser Glu Cys Ser Ala Arg Leu Leu Gln Gln
 1 5 10 15
 Glu Glu Glu Ile Lys Ser Leu Thr Ala Glu Ile Asp Arg Leu Lys
 20 25 30
 Asn Cys Gly Cys Leu Gly Ala Ser Pro Asn Leu Glu Gln Leu Gln
 35 40 45
 Glu Glu Asn Leu Lys Leu Lys Tyr Arg Leu Asn Ile Leu Arg Lys
 50 55 60
 Ser Leu Gln Ala Glu Arg Asn Lys Pro Thr Lys Asn Met Ile Asn
 65 70 75
 Ile Ile Ser Arg Leu Gln Glu Val Phe Gly His Ala Ile Lys Ala
 80 85 90
 Ala Tyr Pro Asp Leu Glu Asn Pro Pro Leu Leu Val Thr Pro Ser
 95 100 105
 Gln Gln Ala Lys Phe Gly Asp Tyr Gln Cys Asn Ser Ala Met Gly
 110 115 120
 Ile Ser Gln Met Leu Lys Thr Lys Glu Gln Lys Val Asn Pro Arg
 125 130 135
 Glu Ile Ala Glu Asn Ile Thr Lys His Leu Pro Asp Asn Glu Cys
 140 145 150
 Ile Glu Lys Val Glu Ile Ala Gly Pro Gly Phe Ile Asn Val His
 155 160 165
 Leu Arg Lys Asp Phe Val Ser Glu Gln Leu Thr Ser Leu Leu Val
 170 175 180
 Asn Gly Val Gln Leu Pro Ala Leu Gly Glu Asn Lys Lys Val Ile
 185 190 195
 Val Asp Phe Ser Ser Pro Asn Ile Ala Lys Glu Met His Val Gly
 200 205 210
 His Leu Arg Ser Thr Ile Ile Gly Glu Ser Ile Ser Arg Leu Phe
 215 220 225
 Glu Phe Ala Gly Tyr Asp Val Leu Arg Leu Asn His Val Gly Asp
 230 235 240
 Trp Gly Thr Gln Phe Gly Met Leu Ile Ala His Leu Gln Asp Lys
 245 250 255
 Phe Pro Asp Tyr Leu Thr Val Ser Pro Pro Ile Gly Asp Leu Gln
 260 265 270
 Val Phe Tyr Lys Glu Ser Lys Lys Arg Phe Asp Thr Glu Glu Glu
 275 280 285
 Phe Lys Lys Arg Ala Tyr Gln Cys Val Val Leu Leu Gln Gly Lys
 290 295 300
 Asn Pro Asp Ile Thr Lys Ala Trp Lys Leu Ile Cys Asp Val Ser
 305 310 315
 Arg Gln Glu Leu Asn Lys Ile Tyr Asp Ala Leu Asp Val Ser Leu
 320 325 330
 Ile Glu Arg Gly Glu Ser Phe Tyr Gln Asp Arg Met Asn Asp Ile
 335 340 345
 Val Lys Glu Phe Glu Asp Arg Gly Phe Val Gln Val Asp Asp Gly
 350 355 360
 Arg Lys Ile Val Phe Val Pro Gly Cys Ser Ile Pro Leu Thr Ile
 365 370 375

Val	Lys	Ser	Asp	Gly	Gly	Tyr	Thr	Tyr	Asp	Thr	Ser	Asp	Leu	Ala	380	385	390
Ala	Ile	Lys	Gln	Arg	Leu	Phe	Glu	Glu	Lys	Ala	Asp	Met	Ile	Ile	395	400	405
Tyr	Val	Val	Asp	Asn	Gly	Gln	Ser	Val	His	Phe	Gln	Thr	Ile	Phe	410	415	420
Ala	Ala	Ala	Gln	Met	Ile	Gly	Trp	Tyr	Asp	Pro	Lys	Val	Thr	Arg	425	430	435
Val	Phe	His	Ala	Gly	Phe	Gly	Val	Val	Leu	Gly	Glu	Asp	Lys	Lys	440	445	450
Lys	Phe	Lys	Thr	Arg	Ser	Gly	Glu	Thr	Val	Arg	Leu	Met	Asp	Leu	455	460	465
Leu	Gly	Glu	Gly	Leu	Lys	Arg	Ser	Met	Asp	Lys	Leu	Lys	Glu	Lys	470	475	480
Glu	Arg	Asp	Lys	Val	Leu	Thr	Ala	Glu	Glu	Leu	Asn	Ala	Ala	Gln	485	490	495
Thr	Ser	Val	Ala	Tyr	Gly	Cys	Ile	Lys	Tyr	Ala	Asp	Leu	Ser	His	500	505	510
Asn	Arg	Leu	Asn	Asp	Tyr	Ile	Phe	Ser	Phe	Asp	Lys	Met	Leu	Asp	515	520	525
Asp	Arg	Gly	Asn	Thr	Ala	Ala	Tyr	Leu	Leu	Tyr	Ala	Phe	Thr	Arg	530	535	540
Ile	Arg	Ser	Ile	Ala	Arg	Leu	Ala	Asn	Ile	Asp	Glu	Glu	Met	Leu	545	550	555
Gln	Lys	Ala	Ala	Arg	Glu	Thr	Lys	Ile	Leu	Leu	Asp	His	Glu	Lys	560	565	570
Glu	Trp	Lys	Leu	Gly	Arg	Cys	Ile	Leu	Arg	Phe	Pro	Glu	Ile	Leu	575	580	585
Gln	Lys	Ile	Leu	Asp	Asp	Leu	Phe	Leu	His	Thr	Leu	Cys	Asp	Tyr	590	595	600
Ile	Tyr	Glu	Leu	Ala	Thr	Ala	Phe	Thr	Glu	Phe	Tyr	Asp	Ser	Cys	605	610	615
Tyr	Cys	Val	Glu	Lys	Asp	Arg	Gln	Thr	Gly	Lys	Ile	Leu	Lys	Val	620	625	630
Asn	Met	Trp	Arg	Met	Leu	Leu	Cys	Glu	Ala	Val	Ala	Ala	Val	Met	635	640	645
Ala	Lys	Gly	Phe	Asp	Ile	Leu	Gly	Ile	Lys	Pro	Val	Gln	Arg	Met	650	655	660

<210> 7

<211> 725

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2662427CD1

<400> 7

Met	Ala	Ala	Ala	Ser	Ala	Val	Ser	Val	Leu	Leu	Val	Ala	Ala	Glu	1	5	10	15
Arg	Asn	Arg	Trp	His	Arg	Leu	Pro	Ser	Leu	Leu	Leu	Pro	Pro	Arg	20	25	30	35
Thr	Trp	Val	Trp	Arg	Gln	Arg	Thr	Met	Lys	Tyr	Thr	Thr	Ala	Thr	40	45	50	55
Gly	Arg	Asn	Ile	Thr	Lys	Val	Leu	Ile	Ala	Asn	Arg	Gly	Glu	Ile	60	65	70	75
Ala	Cys	Arg	Val	Met	Arg	Thr	Ala	Lys	Lys	Leu	Gly	Val	Gln	Thr	80	85	90	95
Val	Ala	Val	Tyr	Ser	Glu	Ala	Asp	Arg	Asn	Ser	Met	His	Val	Asp	100	105	110	115
Met	Ala	Asp	Glu	Ala	Tyr	Ser	Ile	Gly	Pro	Ala	Pro	Ser	Gln	Gln	120	125	130	135
Ser	Tyr	Leu	Ser	Met	Glu	Lys	Ile	Ile	Gln	Val	Ala	Lys	Thr	Ser	140	145	150	155

Ala	Ala	Gln	Ala	Ile	His	Pro	Gly	Cys	Arg	Phe	Leu	Ser	Glu	Asn
				125					130					135
Met	Glu	Phe	Ala	Glu	Leu	Cys	Lys	Gln	Glu	Gly	Ile	Ile	Phe	Ile
				140					145					150
Gly	Pro	Pro	Pro	Ser	Ala	Ile	Arg	Asp	Met	Gly	Ile	Lys	Ser	Thr
				155					160					165
Ser	Lys	Ser	Ile	Met	Ala	Ala	Ala	Gly	Val	Pro	Val	Val	Glu	Gly
				170					175					180
Tyr	His	Gly	Glu	Asp	Gln	Ser	Asp	Gln	Cys	Leu	Lys	Glu	His	Ala
				185					190					195
Arg	Arg	Ile	Gly	Tyr	Pro	Val	Met	Ile	Lys	Ala	Val	Arg	Gly	Gly
				200					205					210
Gly	Gly	Lys	Gly	Met	Arg	Ile	Val	Arg	Ser	Glu	Gln	Glu	Phe	Gln
				215					220					225
Glu	Gln	Leu	Glu	Ser	Ala	Arg	Arg	Glu	Ala	Lys	Lys	Ser	Phe	Asn
				230					235					240
Asp	Asp	Ala	Met	Leu	Ile	Glu	Lys	Phe	Val	Asp	Thr	Pro	Arg	His
				245					250					255
Val	Glu	Val	Gln	Val	Phe	Gly	Asp	His	His	Gly	Asn	Ala	Val	Tyr
				260					265					270
Leu	Phe	Glu	Arg	Asp	Cys	Ser	Val	Gln	Arg	Arg	His	Gln	Lys	Ile
				275					280					285
Ile	Glu	Glu	Ala	Pro	Ala	Pro	Gly	Ile	Lys	Ser	Glu	Val	Arg	Lys
				290					295					300
Lys	Leu	Gly	Glu	Ala	Ala	Val	Arg	Ala	Ala	Lys	Ala	Val	Asn	Tyr
				305					310					315
Val	Gly	Ala	Gly	Thr	Val	Glu	Phe	Ile	Met	Asp	Ser	Lys	His	Asn
				320					325					330
Phe	Cys	Phe	Met	Glu	Met	Asn	Thr	Arg	Leu	Gln	Val	Glu	His	Pro
				335					340					345
Val	Thr	Glu	Met	Ile	Thr	Gly	Thr	Asp	Leu	Val	Glu	Trp	Gln	Leu
				350					355					360
Arg	Ile	Ala	Ala	Gly	Glu	Lys	Ile	Pro	Leu	Ser	Gln	Glu	Glu	Ile
				365					370					375
Thr	Leu	Gln	Gly	His	Ala	Phe	Glu	Ala	Arg	Ile	Tyr	Ala	Glu	Asp
				380					385					390
Pro	Ser	Asn	Asn	Phe	Met	Pro	Val	Ala	Gly	Pro	Leu	Val	His	Leu
				395					400					405
Ser	Thr	Pro	Arg	Ala	Asp	Pro	Ser	Thr	Arg	Ile	Glu	Thr	Gly	Val
				410					415					420
Arg	Gln	Gly	Asp	Glu	Val	Ser	Val	His	Tyr	Asp	Pro	Met	Ile	Ala
				425					430					435
Lys	Leu	Val	Val	Trp	Ala	Ala	Asp	Arg	Gln	Ala	Ala	Leu	Thr	Lys
				440					445					450
Leu	Arg	Tyr	Ser	Leu	Arg	Gln	Tyr	Asn	Ile	Val	Gly	Leu	Pro	Thr
				455					460					465
Asn	Ile	Asp	Phe	Leu	Leu	Asn	Leu	Ser	Gly	His	Pro	Glu	Phe	Glu
				470					475					480
Ala	Gly	Asn	Val	His	Thr	Asp	Phe	Ile	Pro	Gln	His	His	Lys	Gln
				485					490					495
Leu	Leu	Leu	Ser	Arg	Lys	Ala	Ala	Ala	Lys	Glu	Ser	Leu	Cys	Gln
				500					505					510
Ala	Ala	Leu	Gly	Leu	Ile	Leu	Lys	Glu	Lys	Ala	Met	Thr	Asp	Thr
				515					520					525
Phe	Thr	Leu	Gln	Ala	His	Asp	Gln	Phe	Ser	Pro	Phe	Ser	Ser	Ser
				530					535					540
Ser	Gly	Arg	Arg	Leu	Asn	Ile	Ser	Tyr	Thr	Arg	Asn	Met	Thr	Leu
				545					550					555
Lys	Asp	Gly	Lys	Asn	Asn	Val	Ala	Ile	Ala	Val	Thr	Tyr	Asn	His
				560					565					570
Asp	Gly	Ser	Tyr	Ser	Met	Gln	Ile	Glu	Asp	Lys	Thr	Phe	Gln	Val
				575					580					585
Leu	Gly	Asn	Leu	Tyr	Ser	Glu	Gly	Asp	Cys	Thr	Tyr	Leu	Lys	Cys
				590					595					600
Ser	Val	Asn	Gly	Val	Ala	Ser	Lys	Ala	Lys	Leu	Ile	Ile	Leu	Glu

Asn Thr Ile Tyr	605	Gly Ser Ile Glu Ile	610	Asp	615
Ile Pro Val Pro	620	Val Ser Ser Gln Glu Thr	625	Thr	630
Gln Gly Gly Pro	635	Gly Thr Ile Glu Lys Val	640	Val	645
Phe Val Lys Ala	650	Ala Gly Asp Ser Leu Met	655	Met	660
Val Met Ile Ala	665	Thr Ile Lys Ser Pro Lys	670	Lys	675
Asp Gly Thr Val	680	Arg Glu Gly Ala Gln Ala	685	Ala	690
Asn Arg His Thr	695	Glu Glu Glu Glu Ser Asp	700	Asp	705
Lys Arg Glu Ser	710		715		720
	725				

<210> 8

<211> 644

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2844928CD1

<400> 8

Met Ala Val Tyr Val	1	Gly Met Leu Arg Leu	10	Gly Arg Leu Cys Ala	15
Gly Ser Ser Gly Val	5	Ala Ala Leu Ser Arg Ser	25		30
Trp Gln Glu Ala Arg	20	Arg Phe Leu Ser Ser Arg	40		45
Glu Val Asp Arg Met	35	Ile Gly Gly Leu Ser Tyr	55		60
Val Gln Gly Cys Thr	50	Asn Ser Lys Thr Val Gly	70		75
Gln Cys Leu Glu Thr	65	Val Pro Glu Arg Glu Ala	85		90
Leu Val Val Leu His	80	Leu Thr Phe Ala Gln Leu	100		105
Lys Glu Glu Val Asp	95	Gly Leu Leu Ser Ile Gly	115		120
Leu Cys Lys Gly Asp	110	Trp Gly Pro Asn Ser Tyr	130		135
Ala Trp Val Leu Met	125	Ala Gln Ala Gly Ile Ile	145		150
Leu Val Ser Val Asn	140	Ala Met Glu Leu Glu Tyr	160		165
Val Leu Lys Lys Val	155	Leu Val Phe Pro Lys Gln	175		180
Phe Lys Thr Gln Gln	170	Leu Lys Gln Ile Cys Pro	190		195
Glu Val Glu Asn Ala	185	Leu Lys Ser Gln Arg Leu	205		210
Pro Asp Leu Thr Thr	200	Asp Ala Pro Leu Pro Gly	220		225
Thr Leu Leu Leu Asp	215	Ala Gly Ser Thr Arg Gln	235		240
His Leu Asp Gln Leu	230	Gln Phe Leu Ser Cys His	250		255
Asp Pro Ile Asn Ile	245	Gly Thr Thr Gly Ser Pro	265		270
Lys Gly Ala Thr Leu	260	Ile Val Asn Asn Ser Asn	280		285
	275				


```

Ile Leu Gly Glu Arg Leu Lys Leu His Glu Lys Thr Pro Glu Gln
290 295 300
Leu Arg Met Ile Leu Pro Asn Pro Leu Tyr His Cys Leu Gly Ser
305 310 315
Val Ala Gly Thr Met Met Cys Leu Met Tyr Gly Ala Thr Leu Ile
320 325 330
Leu Ala Ser Pro Ile Phe Asn Gly Lys Lys Ala Leu Glu Ala Ile
335 340 345
Ser Arg Glu Arg Gly Thr Phe Leu Tyr Gly Thr Pro Thr Met Phe
350 355 360
Val Asp Ile Leu Asn Gln Pro Asp Phe Ser Ser Tyr Asp Ile Ser
365 370 375
Thr Met Cys Gly Gly Val Ile Ala Gly Ser Pro Ala Pro Pro Glu
380 385 390
Leu Ile Arg Ala Ile Ile Asn Lys Ile Asn Met Lys Asp Leu Val
395 400 405
Val Ala Tyr Gly Thr Thr Glu Asn Ser Pro Val Thr Phe Ala His
410 415 420
Phe Pro Glu Asp Thr Val Glu Gln Lys Ala Glu Ser Val Gly Arg
425 430 435
Ile Met Pro His Thr Glu Ala Arg Ile Met Asn Met Glu Ala Gly
440 445 450
Thr Leu Ala Lys Leu Asn Thr Pro Gly Glu Leu Cys Ile Arg Gly
455 460 465
Tyr Cys Val Met Leu Gly Tyr Trp Gly Glu Pro Gln Lys Thr Glu
470 475 480
Glu Ala Val Asp Gln Asp Lys Trp Tyr Trp Thr Gly Asp Val Ala
485 490 495
Thr Met Asn Glu Gln Gly Phe Cys Lys Ile Val Gly Arg Ser Lys
500 505 510
Asp Met Ile Ile Arg Gly Gly Glu Asn Ile Tyr Pro Ala Glu Leu
515 520 525
Glu Asp Phe Phe His Thr His Pro Lys Val Gln Glu Val Gln Val
530 535 540
Arg His Leu Ala Gln Val Ser Pro Gln Lys Gln Glu Thr His Met
545 550 555
Asn Thr Val Met Ser Asp Ile Phe Leu Trp Pro Trp Asn Val Val
560 565 570
Gly Val Lys Asp Asp Arg Met Gly Glu Glu Ile Cys Ala Cys Ile
575 580 585
Arg Leu Lys Asp Gly Glu Glu Thr Thr Val Glu Glu Ile Lys Ala
590 595 600
Phe Cys Lys Gly Lys Ile Ser His Phe Lys Ile Pro Lys Tyr Ile
605 610 615
Val Phe Val Thr Asn Tyr Pro Leu Thr Ile Ser Gly Lys Ile Gln
620 625 630
Lys Phe Lys Leu Arg Glu Gln Met Glu Arg His Leu Asn Leu
635 640

```

<210> 9
 <211> 504
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 3231586CD1

```

<400> 9
Met Phe Pro Arg Glu Lys Thr Trp Asn Ile Ser Phe Ala Gly Cys
1 5 10 15
Gly Phe Leu Gly Val Tyr Tyr Val Gly Val Ala Ser Cys Leu Arg
20 25 30
Glu His Ala Pro Phe Leu Val Ala Asn Ala Thr His Ile Tyr Gly
35 40 45
Ala Ser Ala Gly Ala Leu Thr Ala Thr Ala Leu Val Thr Gly Val

```

	50		55		60
Cys Leu Gly Glu Ala Gly Ala Lys Phe Ile Glu Val Ser Lys Glu					
	65		70		75
Ala Arg Lys Arg Phe Leu Gly Pro Leu His Pro Ser Phe Asn Leu					
	80		85		90
Val Lys Ile Ile Arg Ser Phe Leu Leu Lys Val Leu Pro Ala Asp					
	95		100		105
Ser His Glu His Ala Ser Gly Arg Leu Gly Ile Ser Leu Thr Arg					
	110		115		120
Val Ser Asp Gly Glu Asn Val Ile Ile Ser His Phe Asn Ser Lys					
	125		130		135
Asp Glu Leu Ile Gln Ala Asn Val Cys Ser Gly Phe Ile Pro Val					
	140		145		150
Tyr Cys Gly Leu Ile Pro Pro Ser Leu Gln Gly Val Arg Tyr Val					
	155		160		165
Asp Gly Gly Ile Ser Asp Asn Leu Pro Leu Tyr Glu Leu Lys Asn					
	170		175		180
Thr Ile Thr Val Ser Pro Phe Ser Gly Glu Ser Asp Ile Cys Pro					
	185		190		195
Gln Asp Ser Ser Thr Asn Ile His Glu Leu Arg Val Thr Asn Thr					
	200		205		210
Ser Ile Gln Phe Asn Leu Arg Asn Leu Tyr Arg Leu Ser Lys Ala					
	215		220		225
Leu Phe Pro Pro Glu Pro Leu Val Leu Arg Glu Met Cys Lys Gln					
	230		235		240
Gly Tyr Arg Asp Gly Leu Arg Phe Leu Gln Arg Asn Gly Leu Leu					
	245		250		255
Asn Arg Pro Asn Pro Leu Leu Ala Leu Pro Pro Ala Arg Pro His					
	260		265		270
Gly Pro Glu Asp Lys Asp Gln Ala Val Glu Ser Ala Gln Ala Glu					
	275		280		285
Asp Tyr Ser Gln Leu Pro Gly Glu Asp His Ile Leu Glu His Leu					
	290		295		300
Pro Ala Arg Leu Asn Glu Ala Leu Leu Glu Ala Cys Val Glu Pro					
	305		310		315
Thr Asp Leu Leu Thr Thr Leu Ser Asn Met Leu Pro Val Arg Leu					
	320		325		330
Ala Thr Ala Met Met Val Pro Tyr Thr Leu Pro Leu Glu Ser Ala					
	335		340		345
Leu Ser Phe Thr Ile Arg Leu Leu Glu Trp Leu Pro Asp Val Pro					
	350		355		360
Glu Asp Ile Arg Trp Met Lys Glu Gln Thr Gly Ser Ile Cys Gln					
	365		370		375
Tyr Leu Val Met Arg Ala Lys Arg Lys Leu Gly Arg His Leu Pro					
	380		385		390
Ser Arg Leu Pro Glu Gln Val Glu Leu Arg Arg Val Gln Ser Leu					
	395		400		405
Pro Ser Val Pro Leu Ser Cys Ala Ala Tyr Arg Glu Ala Leu Pro					
	410		415		420
Gly Trp Met Arg Asn Asn Leu Ser Leu Gly Asp Ala Leu Ala Lys					
	425		430		435
Trp Glu Glu Cys Gln Arg Gln Leu Leu Leu Gly Leu Phe Cys Thr					
	440		445		450
Asn Val Ala Phe Pro Pro Glu Ala Leu Arg Met Arg Ala Pro Ala					
	455		460		465
Asp Pro Ala Pro Ala Pro Ala Asp Pro Ala Ser Pro Gln His Gln					
	470		475		480
Leu Ala Gly Pro Ala Pro Leu Leu Ser Thr Pro Ala Pro Glu Ala					
	485		490		495
Arg Pro Val Ile Gly Ala Leu Gly Leu					
	500				

<210> 10

<211> 489

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3580770CD1

<400> 10

```

Met Lys Tyr Ile Leu Val Thr Gly Gly Val Ile Ser Gly Ile Gly
 1      5      10      15
Lys Gly Ile Ile Ala Ser Ser Ile Gly Thr Ile Leu Lys Ser Cys
 20      25      30
Gly Leu Arg Val Thr Ala Ile Lys Ile Asp Pro Tyr Ile Asn Ile
 35      40      45
Asp Ala Gly Thr Phe Ser Pro Tyr Glu His Gly Glu Val Phe Val
 50      55      60
Leu Asn Asp Gly Gly Glu Val Asp Leu Asp Leu Gly Asp Tyr Glu
 65      70      75
Arg Phe Leu Asp Ile Asn Leu Tyr Lys Asp Thr Ile Val Thr Thr
 80      85      90
Gly Lys Ile Tyr Gln His Val Ile Asn Lys Glu Arg Arg Gly Asp
 95      100      105
Tyr Leu Gly Lys Thr Val Gln Val Val Pro His Ile Thr Asp Ala
 110      115      120
Val Gln Glu Trp Val Met Asn Gln Ala Lys Val Pro Val Asp Gly
 125      130      135
Asn Lys Glu Glu Pro Gln Ile Cys Val Ile Glu Leu Gly Gly Thr
 140      145      150
Ile Gly Asp Ile Glu Gly Met Pro Phe Val Glu Ala Phe Arg Gln
 155      160      165
Phe Gln Phe Lys Ala Lys Arg Glu Asn Phe Cys Asn Ile His Val
 170      175      180
Ser Leu Val Pro Gln Leu Ser Ala Thr Gly Glu Gln Lys Thr Lys
 185      190      195
Pro Thr Gln Asn Ser Val Arg Ala Leu Arg Gly Leu Gly Leu Ser
 200      205      210
Pro Asp Leu Ile Val Cys Arg Ser Ser Thr Pro Ile Glu Met Ala
 215      220      225
Val Lys Glu Lys Ile Ser Met Phe Cys His Val Asn Pro Glu Gln
 230      235      240
Val Ile Cys Ile His Asp Val Ser Ser Thr Tyr Arg Val Pro Val
 245      250      255
Leu Leu Glu Glu Gln Ser Ile Val Lys Tyr Phe Lys Glu Arg Leu
 260      265      270
His Leu Pro Ile Gly Asp Ser Ala Ser Asn Leu Leu Phe Lys Trp
 275      280      285
Arg Asn Met Ala Asp Arg Tyr Glu Arg Leu Gln Lys Ile Cys Ser
 290      295      300
Ile Ala Leu Val Gly Lys Tyr Thr Lys Leu Arg Asp Cys Tyr Ala
 305      310      315
Ser Val Phe Lys Ala Leu Glu His Ser Ala Leu Ala Ile Asn His
 320      325      330
Lys Leu Asn Leu Met Val Ile Asp Met Pro Glu His Asn Pro Gly
 335      340      345
Asn Leu Gly Gly Thr Met Arg Leu Gly Ile Arg Arg Thr Val Phe
 350      355      360
Lys Thr Glu Asn Ser Ile Leu Arg Lys Leu Tyr Gly Asp Val Pro
 365      370      375
Phe Ile Glu Glu Arg His Arg His Arg Phe Glu Val Asn Pro Asn
 380      385      390
Leu Ile Lys Gln Phe Glu Gln Asn Asp Leu Ser Phe Val Gly Gln
 395      400      405
Asp Val Asp Gly Asp Arg Met Glu Ile Ile Glu Leu Ala Asn His
 410      415      420
Pro Tyr Phe Val Gly Val Gln Phe His Pro Glu Phe Ser Ser Arg
 425      430      435
Pro Met Lys Pro Ser Pro Pro Tyr Leu Gly Leu Leu Leu Ala Ala
 440      445      450

```

Thr Gly Asn Leu Asn Ala Tyr Leu Gln Gln Gly Cys Lys Leu Ser
 455 460 465
 Ser Ser Asp Arg Tyr Ser Asp Ala Ser Asp Asp Ser Phe Ser Glu
 470 475 480
 Pro Arg Ile Ala Glu Leu Glu Ile Ser
 485

<210> 11
 <211> 258
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 3778612CD1

<400> 11
 Met Glu Arg Gln Lys Arg Lys Ala Asp Ile Glu Lys Gly Leu Gln
 1 5 10 15
 Phe Ile Gln Ser Thr Leu Pro Leu Lys Gln Glu Glu Tyr Glu Ala
 20 25 30
 Phe Leu Leu Lys Leu Val Gln Asn Leu Phe Ala Glu Gly Asn Asp
 35 40 45
 Leu Phe Arg Glu Lys Asp Tyr Lys Gln Ala Leu Val Gln Tyr Met
 50 55 60
 Glu Gly Leu Asn Val Ala Asp Tyr Ala Ala Ser Asp Gln Val Ala
 65 70 75
 Leu Pro Arg Glu Leu Leu Cys Lys Leu His Val Asn Arg Ala Ala
 80 85 90
 Cys Tyr Phe Thr Met Gly Leu Tyr Glu Lys Ala Leu Glu Asp Ser
 95 100 105
 Glu Lys Ala Leu Gly Pro Asp Ser Glu Ser Ile Arg Ala Leu Phe
 110 115 120
 Arg Lys Ala Arg Ala Leu Asn Glu Leu Gly Arg His Lys Glu Ala
 125 130 135
 Tyr Glu Cys Ser Ser Arg Cys Ser Leu Ala Leu Pro His Asp Glu
 140 145 150
 Ser Val Thr Gln Leu Gly Gln Gly Pro Leu Gly Ser Gly Ala Ser
 155 160 165
 Trp Pro Gly Gln Ser Trp Ser Pro His Arg Val Arg Lys Arg Glu
 170 175 180
 Trp Glu Ala Glu Cys Asp Gly Glu Glu Gly Gln Glu Asp Pro Phe
 185 190 195
 Asn Asp Glu Gly Asn Tyr Phe Ser Cys Glu Pro Ser Arg Ala Pro
 200 205 210
 Gly Trp Glu Ala Gln Arg Thr Glu Ser Gly Thr Cys Val Pro Pro
 215 220 225
 Gly Arg Gln Gly Gln Asp Gly Met Ala Ser Met Gly Ala Gly Trp
 230 235 240
 Val Gly Arg Asp Ala Ala Phe Leu Ser Trp Ala Val Ile Asn Leu
 245 250 255
 Met Val Leu

<210> 12
 <211> 555
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 4574912CD1

<220>
 <221> unsure
 <222> 63-65
 <223> unknown or other

<400> 12

Met	Glu	Gly	Ala	Val	Leu	Glu	Ala	Gly	Gly	Ala	Arg	Cys	Phe	Cys
1				5					10					15
Arg	Phe	Gly	Cys	Glu	Leu	Ser	Lys	Tyr	Glu	Asn	Pro	Gly	Tyr	Ser
				20					25					30
Ser	Pro	Arg	Ser	Asp	Tyr	Phe	Lys	Asn	Tyr	Met	Ile	Ile	Ile	Thr
				35					40					45
Gln	Asn	Arg	Met	Ser	Phe	Leu	Ala	Asn	Met	Phe	His	Thr	Met	Asp
				50					55					60
Cys	Val	Xaa	Xaa	Xaa	Arg	Tyr	Ser	Cys	Gly	Pro	Thr	Val	Tyr	Asp
				65					70					75
His	Ala	His	Leu	Gly	His	Ala	Cys	Ser	Tyr	Val	Arg	Phe	Asp	Ile
				80					85					90
Ile	Arg	Arg	Ile	Leu	Thr	Lys	Val	Phe	Gly	Cys	Ser	Ile	Val	Met
				95					100					105
Val	Met	Gly	Ile	Thr	Asp	Val	Asp	Asp	Lys	Ile	Ile	Lys	Arg	Ala
				110					115					120
Asn	Glu	Met	Asn	Ile	Ser	Pro	Ala	Ser	Leu	Ala	Ser	Leu	Tyr	Glu
				125					130					135
Glu	Asp	Phe	Lys	Gln	Asp	Met	Ala	Ala	Leu	Lys	Val	Leu	Pro	Pro
				140					145					150
Thr	Val	Tyr	Leu	Arg	Val	Thr	Glu	Asn	Ile	Pro	Gln	Ile	Ile	Ser
				155					160					165
Phe	Ile	Glu	Gly	Ile	Ile	Ala	Arg	Gly	Asn	Ala	Tyr	Ser	Thr	Ala
				170					175					180
Lys	Gly	Asn	Val	Tyr	Phe	Asp	Leu	Lys	Ser	Arg	Gly	Asp	Lys	Tyr
				185					190					195
Gly	Lys	Leu	Val	Gly	Val	Val	Pro	Gly	Pro	Val	Gly	Glu	Pro	Ala
				200					205					210
Asp	Ser	Asp	Lys	Arg	His	Ala	Ser	Asp	Phe	Ala	Leu	Trp	Lys	Ala
				215					220					225
Ala	Lys	Pro	Gln	Glu	Val	Phe	Trp	Ala	Ser	Pro	Trp	Gly	Pro	Gly
				230					235					240
Arg	Pro	Gly	Trp	His	Ile	Glu	Cys	Ser	Ala	Ile	Ala	Ser	Met	Val
				245					250					255
Phe	Gly	Ser	Gln	Leu	Asp	Ile	His	Ser	Gly	Gly	Ile	Asp	Leu	Ala
				260					265					270
Phe	Pro	His	His	Glu	Asn	Glu	Ile	Ala	Gln	Cys	Glu	Val	Phe	His
				275					280					285
Gln	Cys	Glu	Gln	Trp	Gly	Asn	Tyr	Phe	Leu	His	Ser	Gly	His	Leu
				290					295					300
His	Ala	Lys	Gly	Lys	Glu	Glu	Lys	Met	Ser	Lys	Ser	Leu	Lys	Asn
				305					310					315
Tyr	Ile	Thr	Ile	Lys	Asp	Phe	Leu	Lys	Thr	Phe	Ser	Pro	Asp	Val
				320					325					330
Phe	Arg	Phe	Phe	Cys	Leu	Arg	Ser	Ser	Tyr	Arg	Ser	Ala	Ile	Asp
				335					340					345
Tyr	Ser	Asp	Ser	Ala	Met	Leu	Gln	Ala	Gln	Gln	Leu	Leu	Leu	Gly
				350					355					360
Leu	Gly	Ser	Phe	Leu	Glu	Asp	Ala	Arg	Ala	Tyr	Met	Lys	Gly	Gln
				365					370					375
Leu	Ala	Cys	Gly	Ser	Val	Arg	Glu	Ala	Met	Leu	Trp	Glu	Arg	Leu
				380					385					390
Ser	Ser	Thr	Lys	Arg	Ala	Val	Lys	Ala	Ala	Leu	Ala	Asp	Asp	Phe
				395					400					405
Asp	Thr	Pro	Arg	Val	Val	Asp	Ala	Ile	Leu	Gly	Leu	Ala	His	His
				410					415					420
Gly	Asn	Gly	Gln	Leu	Arg	Ala	Ser	Leu	Lys	Glu	Pro	Glu	Gly	Pro
				425					430					435
Arg	Ser	Pro	Ala	Val	Phe	Gly	Ala	Ile	Ile	Ser	Tyr	Phe	Glu	Gln
				440					445					450
Phe	Phe	Glu	Thr	Val	Gly	Ile	Ser	Leu	Ala	Asn	Gln	Gln	Tyr	Val
				455					460					465
Ser	Gly	Asp	Gly	Ser	Glu	Ala	Thr	Leu	His	Gly	Val	Val	Asp	Glu
				470					475					480

Leu	Val	Arg	Phe	Arg	Gln	Lys	Val	Arg	Gln	Phe	Ala	Leu	Ala	Met
				485					490					495
Pro	Glu	Ala	Thr	Gly	Asp	Ala	Arg	Arg	Gln	Gln	Leu	Leu	Glu	Arg
				500					505					510
Gln	Pro	Leu	Leu	Glu	Ala	Cys	Asp	Thr	Leu	Arg	Arg	Gly	Leu	Thr
				515					520					525
Ala	His	Gly	Ile	Asn	Ile	Lys	Asp	Arg	Ser	Ser	Thr	Thr	Ser	Thr
				530					535					540
Trp	Glu	Leu	Leu	Asp	Gln	Arg	Thr	Lys	Asp	Gln	Lys	Ser	Ala	Gly
				545					550					555

<210> 13
 <211> 463
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 5630806CD1

<400> 13

Met	Ala	Ala	Ser	Met	Phe	Tyr	Gly	Arg	Leu	Val	Ala	Val	Ala	Thr
1				5					10					15
Leu	Arg	Asn	His	Arg	Pro	Arg	Thr	Ala	Gln	Arg	Ala	Ala	Ala	Gln
				20					25					30
Val	Leu	Gly	Ser	Ser	Gly	Leu	Phe	Asn	Asn	His	Gly	Leu	Gln	Val
				35					40					45
Gln	Gln	Gln	Gln	Gln	Arg	Asn	Leu	Ser	Leu	His	Glu	Tyr	Met	Ser
				50					55					60
Met	Glu	Leu	Leu	Gln	Glu	Ala	Gly	Val	Ser	Val	Pro	Lys	Gly	Tyr
				65					70					75
Val	Ala	Lys	Ser	Pro	Asp	Glu	Ala	Tyr	Ala	Ile	Ala	Lys	Lys	Leu
				80					85					90
Gly	Ser	Lys	Asp	Val	Val	Ile	Lys	Ala	Gln	Val	Leu	Ala	Gly	Gly
				95					100					105
Arg	Gly	Lys	Gly	Thr	Phe	Glu	Ser	Gly	Leu	Lys	Gly	Gly	Val	Lys
				110					115					120
Ile	Val	Phe	Ser	Pro	Glu	Glu	Ala	Lys	Ala	Val	Ser	Ser	Gln	Met
				125					130					135
Ile	Gly	Lys	Lys	Leu	Phe	Thr	Lys	Gln	Thr	Gly	Glu	Lys	Gly	Arg
				140					145					150
Ile	Cys	Asn	Gln	Val	Leu	Val	Cys	Glu	Arg	Lys	Tyr	Pro	Arg	Arg
				155					160					165
Glu	Tyr	Tyr	Phe	Ala	Ile	Thr	Met	Glu	Arg	Ser	Phe	Gln	Gly	Pro
				170					175					180
Val	Leu	Ile	Gly	Ser	Ser	His	Gly	Gly	Val	Asn	Ile	Glu	Asp	Val
				185					190					195
Ala	Ala	Glu	Thr	Pro	Glu	Ala	Ile	Ile	Lys	Glu	Pro	Ile	Asp	Ile
				200					205					210
Glu	Glu	Gly	Ile	Lys	Lys	Glu	Gln	Ala	Leu	Gln	Leu	Ala	Gln	Lys
				215					220					225
Met	Gly	Phe	Pro	Pro	Asn	Ile	Val	Glu	Ser	Ala	Ala	Glu	Asn	Met
				230					235					240
Val	Lys	Leu	Tyr	Ser	Leu	Phe	Leu	Lys	Tyr	Asp	Ala	Thr	Met	Ile
				245					250					255
Glu	Ile	Asn	Pro	Met	Val	Glu	Asp	Ser	Asp	Gly	Ala	Val	Leu	Cys
				260					265					270
Met	Asp	Ala	Lys	Ile	Asn	Phe	Asp	Ser	Asn	Ser	Ala	Tyr	Arg	Gln
				275					280					285
Lys	Lys	Ile	Phe	Asp	Leu	Gln	Asp	Trp	Thr	Gln	Glu	Asp	Glu	Arg
				290					295					300
Asp	Lys	Asp	Ala	Ala	Lys	Ala	Asn	Leu	Asn	Tyr	Ile	Gly	Leu	Asp
				305					310					315
Gly	Asn	Ile	Gly	Cys	Leu	Val	Asn	Gly	Ala	Gly	Leu	Ala	Met	Ala
				320					325					330

```

Thr Met Asp Ile Ile Lys Leu His Gly Gly Thr Pro Ala Asn Phe
      335      340      345
Leu Asp Val Gly Gly Gly Ala Thr Val His Gln Val Thr Glu Ala
      350      355      360
Phe Lys Leu Ile Thr Ser Asp Lys Lys Val Leu Ala Ile Leu Val
      365      370      375
Asn Ile Phe Gly Gly Ile Met Arg Cys Asp Val Ile Ala Gln Gly
      380      385      390
Ile Val Met Ala Val Lys Asp Leu Glu Ile Lys Ile Pro Val Val
      395      400      405
Val Arg Leu Gln Gly Thr Arg Val Asp Asp Ala Lys Ala Leu Ile
      410      415      420
Ala Asp Ser Gly Leu Lys Ile Leu Ala Cys Asp Asp Leu Asp Glu
      425      430      435
Ala Ala Arg Met Val Val Lys Leu Ser Glu Ile Val Thr Leu Ala
      440      445      450
Lys Gln Ala His Val Asp Val Lys Phe Gln Leu Pro Ile
      455      460

```

<210> 14

<211> 399

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5854855CD1

<400> 14

```

Met Tyr Phe Gly Ala Val Ser Trp Leu Cys Gly Pro Arg Ala Pro
  1      5      10      15
Asp Glu Val Ser Arg Arg Pro Asp Pro Arg Lys Gly Gln Leu Gly
      20      25      30
Val Ala Phe Val Leu Leu Pro Pro His Ser Glu Gly Ala Arg Val
      35      40      45
Phe Gly Ala Leu Gly Pro Ile Gly Pro Ser Ser Pro Gly Leu Thr
      50      55      60
Leu Gly Gly Leu Ala Val Ser Glu His Arg Leu Ser Asn Lys Leu
      65      70      75
Leu Ala Trp Ser Gly Val Leu Glu Trp Gln Glu Lys Arg Arg Pro
      80      85      90
Tyr Ser Asp Ser Thr Ala Lys Leu Lys Arg Thr Leu Pro Cys Gln
      95      100      105
Ala Tyr Val Asn Gln Gly Glu Asn Leu Glu Thr Asp Gln Trp Pro
      110      115      120
Gln Lys Leu Ile Met Gln Leu Ile Pro Gln Gln Leu Leu Thr Thr
      125      130      135
Leu Gly Pro Leu Phe Arg Asn Ser Gln Leu Ala Gln Phe His Phe
      140      145      150
Thr Asn Arg Asp Cys Asp Ser Leu Lys Gly Leu Cys Arg Ile Met
      155      160      165
Gly Asn Gly Phe Ala Gly Cys Met Leu Phe Pro His Ile Ser Pro
      170      175      180
Cys Glu Val Arg Val Leu Met Leu Leu Tyr Ser Ser Lys Lys Lys
      185      190      195
Ile Phe Met Gly Leu Ile Pro Tyr Asp Gln Ser Gly Phe Val Ser
      200      205      210
Ala Ile Arg Gln Val Ile Thr Thr Arg Lys Gln Ala Val Gly Pro
      215      220      225
Gly Gly Val Asn Ser Gly Pro Val Gln Ile Val Asn Asn Lys Phe
      230      235      240
Leu Ala Trp Ser Gly Val Met Glu Trp Gln Glu Pro Arg Pro Glu
      245      250      255
Pro Asn Ser Arg Ser Lys Arg Trp Leu Pro Ser His Val Tyr Val
      260      265      270
Asn Gln Gly Glu Ile Leu Arg Thr Glu Gln Trp Pro Arg Lys Leu

```

	275		280		285
Tyr Met Gln Leu Ile	Pro Gln Gln Leu	Leu Thr Thr Leu Val	Pro		
	290		295		300
Leu Phe Arg Asn Ser	Arg Leu Val Gln	Phe His Phe Thr Lys	Asp		
	305		310		315
Leu Glu Thr Leu Lys	Ser Leu Cys Arg	Ile Met Asp Asn Gly	Phe		
	320		325		330
Ala Gly Cys Val His	Phe Ser Tyr Lys	Ala Ser Cys Glu Ile	Arg		
	335		340		345
Val Leu Met Leu Leu	Tyr Ser Ser Glu	Lys Lys Ile Phe Ile	Gly		
	350		355		360
Leu Ile Pro His Asp	Gln Gly Asn Phe	Val Asn Gly Ile Arg	Arg		
	365		370		375
Val Ile Ala Asn Gln	Gln Gln Val Leu	Gln Arg Asn Leu Glu	Gln		
	380		385		390
Glu Gln Gln Gln Arg	Gly Met Gly Gly				
	395				

<210> 15

<211> 339

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5993973CD1

<400> 15

Met Val Ser Gly Cys	Gln Thr Arg Ser	Ile Leu Glu Tyr Leu	Arg		
1	5	10	15		
Val Gly Gly Arg Gly	Gly Gly Lys Gly	Lys Gly Arg Ala Glu	Gly		
	20	25	30		
Ser Glu Lys Glu Glu	Ser Arg Arg Lys	Arg Arg Glu Arg Lys	Gln		
	35	40	45		
Arg Arg Glu Gly Gly	Asp Gly Glu Glu	Gln Asp Val Gly Asp	Ala		
	50	55	60		
Gly Arg Leu Leu Leu	Arg Val Leu His	Val Ser Glu Asn Pro	Val		
	65	70	75		
Pro Leu Thr Val Arg	Val Ser Pro Glu	Val Arg Asp Val Arg	Pro		
	80	85	90		
Tyr Ile Val Gly Ala	Val Val Arg Gly	Met Asp Leu Gln Pro	Gly		
	95	100	105		
Asn Ala Leu Lys Arg	Phe Leu Thr Ser	Gln Thr Lys Leu His	Glu		
	110	115	120		
Asp Leu Cys Glu Lys	Arg Thr Ala Ala	Thr Leu Ala Thr His	Glu		
	125	130	135		
Leu Arg Ala Val Lys	Gly Pro Leu Leu	Tyr Cys Ala Arg Pro	Pro		
	140	145	150		
Gln Asp Leu Lys Ile	Val Pro Leu Gly	Arg Lys Glu Ala Lys	Ala		
	155	160	165		
Lys Glu Leu Val Arg	Gln Leu Gln Leu	Glu Ala Glu Glu Gln	Arg		
	170	175	180		
Lys Gln Lys Lys Arg	Gln Ser Val Ser	Gly Leu His Arg Tyr	Leu		
	185	190	195		
His Leu Leu Asp Gly	Asn Glu Asn Tyr	Pro Cys Leu Val Asp	Ala		
	200	205	210		
Asp Gly Asp Val Ile	Ser Phe Pro Pro	Ile Thr Asn Ser Glu	Lys		
	215	220	225		
Thr Lys Val Lys Lys	Thr Thr Ser Asp	Leu Phe Leu Glu Val	Thr		
	230	235	240		
Ser Ala Thr Ser Leu	Gln Ile Cys Lys	Asp Val Met Asp Ala	Leu		
	245	250	255		
Ile Leu Lys Met Ala	Glu Met Lys Lys	Tyr Thr Leu Glu Asn	Lys		
	260	265	270		
Glu Glu Gly Ser Leu	Ser Asp Thr Glu	Ala Asp Ala Val Ser	Gly		
	275	280	285		

Gln	Leu	Pro	Asp	Pro	Thr	Thr	Asn	Pro	Ser	Ala	Gly	Lys	Asp	Gly
				290					295					300
Pro	Ser	Leu	Leu	Val	Val	Glu	Gln	Val	Arg	Val	Val	Asp	Leu	Glu
				305					310					315
Gly	Ser	Leu	Lys	Val	Val	Tyr	Pro	Ser	Lys	Ala	Asp	Leu	Ala	Thr
				320					325					330
Ala	Pro	Pro	His	Val	Thr	Val	Val	Arg						
				335										

<210> 16
 <211> 3902
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1806212CB1

<400> 16
 ggatgattgc ctcagcaggt gtgaagcgtg tgcttttagtt tcgtgggagg cctggcatcc 60
 ccgagagggg ggggaaaggt aaccactcct ttgtggaggt cgccagggtc attgtcgtgg 120
 atttgcacag tcggctgggc ggtgcaatgg cggaaaagaaa aggaacagcc aaagtggact 180
 ttttgaagaa gattgagaaa gaaatccaac agaaatggga tactgagaga gtgtttgagg 240
 tcaatgcac taaatttagag aaacagacca gcaagggcaa gtattttgta accttcccat 300
 acccatatat gaatggacgc cttcatttgg gacacacgtt ttctttatcc aaatgtgagt 360
 ttgctgtagg gtaccagcga ttgaaaggaa aatgttgtct gtttcccttt ggcctgcact 420
 gtactggaat gcctattaag gcatgtgctg ataagttgaa aagagaaaata gagctgtatg 480
 gttgcccccc tgattttcca gatgaagaag aggaagagga agaaaccagt gttaaaacag 540
 aagatataat aattaaggat aaagctaaag gaaaaaagag taaagctgct gctaaagctg 600
 gatcttctaa ataccagtg ggcattatga aatcccttgg cctgtctgat gaagagatag 660
 taaaattttc tgaagcagaa cattggcctt attatttccc gccactggct attcaggatt 720
 taaaaagaat ggggttgaag gtagactggc gtcgttcctt catcaccact gatgttaatc 780
 cttactatga ttcatattgtc agatggcaat ttttaacatt aagagaaaaga aacaaaatta 840
 aatttgggaa gcggtatata atttactctc cgaaagatgg acagccttgc atggatcatg 900
 atagacaaac tggagagggt gttggacctc aggaatatac tttactcaaa ttgaaggtgc 960
 ttgagccata cccatctaaa ttaagtggcc tgaaaggtaa aaatatattc ttggtggctg 1020
 ctactctcag acctgagacc atgtttgggc agacaaattg ttgggttcgt cctgatatga 1080
 agtacattgg atttgagacg gtgaatggtg atatatcat ctgtacccaa aaagcagcca 1140
 ggaatatgtc ataccagggc ttaccaaaag acaatggcgt ggtgcctgtt gttaaggaaat 1200
 taatggggga ggaatttctt ggtgcacac tttctgcacc ttttaacatca tacaaggtga 1260
 tctatgttct cccaatgcta actattaagg aggataaagg cactgggtgtg gttacaagtg 1320
 ttcttccga ctccctgat gatattgctg cctcagaga cttgaagaaa aagcaagcct 1380
 tacgagcaaa atatggaatt agagatgaca tgggtcttgc atttgagccg gtgccagtca 1440
 ttgaaatccc aggttttggg aatctttctg ctgtaaccat ttgtgatgag ttgaaaattc 1500
 agagccagaa tgaccgggaa aaacttgcag aagcaaaagg gaagatatat ctaaaaggat 1560
 tttatgaggy tatcatgttg gtggatggat ttaaaggaca gaagggttcaa gatgtaaaga 1620
 agactattca gaaaaagatg attgacgctg gagatgcact tatttacatg gaaccagaga 1680
 aacaagtgat gtccaggtcg tcagatgaat gtgttgtggc tctgtgtgac cagtggtagt 1740
 tggattatgg agaagagaat tggaagaaac agacatctca gtgcttgaag aacctggaaa 1800
 cattctgtga ggagaccagg aggaattttg aagccacctt aggttggcta caagaacatg 1860
 cttgctcaag aacttatggg ctaggcactc acctgccttg ggatgagcag tggctgattg 1920
 aatcactttc tgactccact atttacatgg cattttacac agttgcacac ctattgcagg 1980
 ggggtaactt gcatggacag gcagagtctc cgctgggcat tagaccgcaa cagatgacca 2040
 aggaagtttg ggattatgtt ttcttcaagg aggtccatt tcctaagact cagattgcaa 2100
 aggaaaaaatt agatcagtta aagcaggagt ttgaattctg gtatcctgtt gatcttcgag 2160
 tctctggcaa ggatcttgtt ccaaatacat tttcatatta cctttataat catgtggcta 2220
 tgtggccgga acaaagtgc aaatggccta cagctgtgag agcaaatgga catctcctcc 2280
 tgaactctga gaagatgtca aaatccacag gcaacttctc cactttgacc caagctattg 2340
 acaaattttc agcagatgga atgcgttttg ctctggctga tgctgggtgac actgtagaag 2400
 atgccaaact tgtggaagcc atggcagatg caggtattct ccgtctgtac acctgggtag 2460
 agtgggtgaa agaaatggtt gccaactggg acagcctaag aagtggctct gccagcactt 2520
 tcaatgatag agtttttggc agtgaattga atgcaggaa tataaaaaaca gatcaaaact 2580
 atgaaaagat gatgttttaa gaagctttga aaacagggtt ttttgagttt caggccgcaa 2640
 aagataagta ccgtgaattg gctgtggaag gtagtcacag agaacttgtg ttccggttta 2700
 ttgaagtcca gacacttctc ctctgtccat tctgtccaca tttgtgtgag cacatctgga 2760
 cactcctggg aaagcctgac tcaattatga atgttcatg gcctgtggca ggtcctgtta 2820

```

atgaagtttt aatacactcc tcacagtatc ttatggaagt aacacatgac cttagactac 2880
gactcaagaa ctatatgatg ccagctaaag ggaagaagac tgacaaacaa cccctgcaga 2940
agccctcaca ttgcaccatc tatgtggcaa agaactatcc accttggcaa cataccaccc 3000
tgtctgttct acgtaaacac tttgaggcca ataacggaaa actgcctgac aacaaagtca 3060
ttgttagtga actaggcagt atgccagaac tgaagaaata catgaagaaa gtcattgccat 3120
ttgttgccat gattaaggaa aatctggaga agatggggcc tcgtattctg gatttgcaat 3180
tagaatttga tgaagggtgt gtgcttatgg agaataatag ctatctgact aattcgcttg 3240
agctagaaca catagaagtc aagtttgcct ccgaagcaga agataaaatc aggggaagact 3300
gctgtcctgg gaaaccactt aatgttttta gaatagaacc tgggtgtgtc gtttctctgg 3360
tgaatcccca gccatccaat ggccacttct caaccaaata tgaaatcagg caaggagata 3420
actgtgatcc cataatcagg cgtttaatga aatgaatcg aggaattaaa gacctttcca 3480
aagtgaact gatgagattt gatgatccac tgttggggcc tcgacgagtt cctgtcctgg 3540
gaaaggagta caccgagaag acccccattt ctgagcatgc tgttttcaat gtggacctca 3600
tgagcaagaa aattcatctg actgagaatg ggataagggt ggataattggc gatacaataa 3660
tctatctggg tcattaaact catgcacatt ggagatttat cctggtttct taggaatact 3720
actactctga ttgtgtctac tgattggcta tcagaacctt aggctggacc taaatagatt 3780
gatttcattt ctaaccatcc aattctgcat gtattcataa ttctatcaag tcatctttga 3840
ttcctggacc taataaattt tttttccctg ttctttgggt gtccaagaaa aaaaaaaaaa 3900
aa

```

<210> 17

<211> 3317

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2083883CB1

<400> 17

```

catggcctga ctcgggacag ctcagagcag ggcagaactg gggacactct gggccggcct 60
tctgcctgca tggacgctct gaagccaccc tgtctctgga ggaaccacga gcgagggaag 120
aaggacaggg actcgtgtgg caggaagaac tcagagccgg gaagcccca ttactagaa 180
gcactgagag atgcggcccc ctcgcagggt ctgaatttcc tgcgtctgtt cacaaagatg 240
ctttttatct ttaacttttt gttttcccca ctccgaccc cggcggtgat ctgcatcctg 300
acatttggag ctgccatctt ctgtggctg atcaccagac ctcaaccctt cttacctctt 360
cttgacctga acaatcagtc tgtgggaatt gagggaggag cacggaaggg ggtttcccag 420
aagaacaatg acctaaacag ttgctgcttc tcagatgcca agactatgta tgaggttttc 480
caaagaggac tcgctgtgtc tgacaatggg ccctgcttgg gatatagaaa accaaaccag 540
ccctacagat ggctatctta caaacagggt tctgatagag cagagtacct gggttcctgt 600
ctcttgcata aaggttataa atcatcaca gaccagtttg tcggcatctt tgctcagaat 660
aggccagagt ggatcatctc cgaattggct tgttacacgt actctatggt agctgtacct 720
ctgtatgaca ccttgggacc agaagccatc gtacatattg tcaacaaggc tgatatcgcc 780
tggtgtatct gtgacacacc ccaaaaggca ttggtgctga tagggaatgt agagaaaggg 840
ttcaccccgga gctgaaggt gatcctcctt atggaccctt ttgatgatga cctgaagcaa 900
agaggggaga agagtggaaat tgagatctta tccctatatg atgctgagaa cctaggcaaa 960
gagcacttca gaaaacctgt gcctcctagc ccagaagacc tgagcgtcat ctgcttcacc 1020
agtgggacca caggtgaccc caaaggagcc atgataaccc atcaaaatat tgtttcaaat 1080
gctgctgctt ttctcaaatg tgtggagcat gcttatgagc ccactcctga tgatgtggcc 1140
atatactacc tccctctggc tcatatgttt gagaggattg tacaggctgt tgtgtacagc 1200
tgtggagcca gagttggatt cttccaaggg gatattcggg tgctggctga cgacatgaag 1260
actttgaagc ccacattgtt tcccgcggtg cctcgactcc ttaacaggat ctacgataag 1320
gtacaaaatg aggccaaagc acccttgaag aagttcttgt tgaagctggc tgtttccagt 1380
aaattcaaaag agcttcaaaa gggatatcat aggcattgata gtttctggga caagctcatc 1440
tttgcaaaga tccaggacag cctgggcgga agggttcgtg taattgtcac tggagctgcc 1500
cccatgtcca cttcagtcac gacattcttc cgggcagcaa tgggatgtca ggtgtatgaa 1560
gcttatggtc aaacagaatg cacagggtggc tgtacattta cattacctgg ggactggaca 1620
tcaggctcac ttggggctgc cctggcttgc aattacgtga agctggaaga tgtggctgac 1680
atgaactact ttacagtga taatgaagga gaggtctgca tcaagggtag aaacgtgttc 1740
aaaggatacc tgaaggaccc tgagaagaca caggaagccc tggacagtga tggctggctt 1800
cacacaggag acattggctg ctggctcccc aatggaactc tgaagatcat cgaccgtaaa 1860
aagaacattt tcaagctggc ccaaggagaa tatattgcac cagagaagat agaaaaatc 1920
tcaacagga gtcaaccagt gttacaaatt ttgtacacg gggagagctt acggctatcc 1980
ttagtaggag tgggtggttc tgacacagat gtaactccct catttgcagc caagcttggg 2040
gtgaagggtc cctttgagga actgtgccaa aaccaagttg taagggaagc cattttagaa 2100

```

```

gacttgcaga aaattgggaa agaaagtggc cttaaaactt ttgaacaggt caaagccatt 2160
tttcttcatc cagagccatt ttccattgaa aatgggctct tgacaccaac attgaaagca 2220
aagcgaggag agctttccaa atactttcgg acccaaatg acagcctgta tgagcacatc 2280
caggattagg ataaggtact taagtacctg ccggccact gtgcactgct tgtgagaaaa 2340
tggattaaaa actattctta catttgtttt gcctttcctc ctattttttt ttaacctgtt 2400
aaactctaaa gccatagctt ttgttttata ttgagacata taatgtgtaa acttagttcc 2460
caaataaatc aatcctgtct ttcccatctt cgatgttgct aatattaagg cttcaggggt 2520
acttttatca acatgcctgt cttcaagatc ccagtttatg ttctgtgtcc ttcctcatga 2580
tttccaacct taatactatt agtaaccaca agttcaaggg tcaaaggac cctctgtgcc 2640
ttcttctttg ttttgtgata aacataactt gccaacagtc tctatgctta tttacatctt 2700
ctactgttca aactaagaga tttttaaatt ctgaaaaact gcttacaatt catgttttct 2760
agccactcca caaaccacta aaattttagt tttagcctat cactcatgtc aatcatatct 2820
atgagacaaa tgtctccgat gctcttctgc gtaaatataa ttgtgtactg aagggaaaag 2880
tttgatcata ccaaacattt cctaaactct ctagttagat atctgacttg ggagtattaa 2940
aaattgggtc tatgacatac tgtccaaaag gaatgctgtt cttaaagcat tatttacagt 3000
aggaactggg gagtaaatct gtccctaca gtttgctgct gagctggaag ctgtggggga 3060
aggagtgtac aggtggggcc agtgaacttt tccagtaaat gaagcaagca ctgaataaaa 3120
acctcctgaa ctgggaacaa agatctacag gcaagcaaga tgccacaca acaggcttat 3180
tttctgtgaa ggaaccaact gatctcccc acccttggat tagagtctct gctctacctt 3240
acccacagat aacacatgtt gtttctactt gtaaatgtaa agtctttaa ataaactatt 3300
acagataaaa aaaaaaa 3317

```

<210> 18

<211> 1928

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2454288CB1

<400> 18

```

cccacgcgtc cgcgacacac catgccgact gtcagcgtga agcgtgatct gctcttcaa 60
gccctgggcc gcacctacac tgacgaagaa tttgatgaac tatgttttga atttggctg 120
gagcttgatg aaattacatc tgagaaggaa ataataagta aagaacaagg taatgtaaag 180
gcagcaggag cctctgatgt tgttctttac aaaattgacg tcctggccaa tagatatgat 240
ctcctgtgtc tggaaggatt ggttcgagga cttcaggtct tcaaagaaag gataaaggct 300
ccagtgtata aacgggtaat gcctgatgga aaaatccaga aattgattat cacagaagag 360
acagctaaga tacgtccttt tgccggtagca gcagttctcc gtaataataa gtttactaaa 420
gatcgatatg acagcttcat tgaacttcag gagaaattac atcagaatat ttgcaggaaa 480
agagcactgg ttgccattgg tacccatgat ttggacactt tgtcgggccc atttacttat 540
actgcaaagc gtccttcaga tatcaaattc aagcctctaa ataagaccaa ggagtataca 600
gcctgtgaac tgatgaacat atacaagact gacaatcacc tgaaacatta tttacatc 660
attgaaaaca aaccctgtga tccagttatc tatgatagca atggtgtcgt cctttcaatg 720
cctcccatca tcaatgggga tcattccaga ataacagtaa atactagaaa tatttttatt 780
gaatgcacgg gaactgactt tactaaggca aaaatagttc ttgatattat tgtcaccatg 840
ttcagtgaat attgtgagaa tcaatttacg gtcgaagctg ctgaagtggg ttttcctaat 900
ggaaaatcac atacctttcc agaattagct taccgaaagg agatggtag agctgacct 960
attaacaaaa aagttggaat cagagaaact ccagaaaatc ttgccaaact tctgaccagg 1020
atgtatttaa aatcagaagt cataggtgat gggaaatcaga ttgagattga aatccctcca 1080
accagagctg acattatcca tgcagtgtat attgtagaag atgcagctat tgcttatgga 1140
tataacaaca ttcagatgac tctcccgaaa acttacacca tagctaatac atttctctt 1200
aataagctca ctgaacttct ccgacatgac atggcagccg ctggcttcac tgaagcactt 1260
acctttgccc tgtgtctcca agaagatatt gctgataaac taggtgtgga tatctctgca 1320
acaaaggcag tccacataag taatcctaaa acagctgaat ttcaggtggc acgcactacc 1380
cttcttctct gctcctgaa gaccatagca gcaaatcgta agatgccct tccactgaaa 1440
ctgtttgaaa tctctgacat tgaataaaa gattctaata cagatgtagg tgcaaaaaac 1500
tacagacatc tctgtctgtt ttattacaac aagaatcctg ggtttgagat cattcatggg 1560
ctgctggaca gaattatgca gttgtctgat gtgctcctg gtgaagacaa ggggggatat 1620
gtgatcaaag catcagaagg gcctgcttct tccccgggc gatgtgcaga gatctttgcc 1680
aggggtcaaa gcgtcgggaa gcttggggtc cttcatcctg acgttatcac caaatttgag 1740
ctgacacatg cctgtcctc ctagaaaatc aatattggac cctttttgtg aagattgggtc 1800
tctgtgtgtt gattctcttc ccaggtgtcc ctttctcctc ccctagtgtc ctttaagtct 1860
cctccacagg gaacatctat ttgggctttg atgtttaata aagtagaaag cactgtcaaa 1920
aaaaaaaaa 1928

```

<210> 19
<211> 2122
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 1513539CB1

<400> 19
cagggtcacac gcagccagtc agcattagag ctgagatgcc aagctctcag ccacacatgt 60
gcactcacct ggaacatctg tggagctgtt agaatecgtg ccacttttca tgaaactgaa 120
gtgcatcttt gggtttgcaa ctaaagaaac cagctgctac aatgtcacta acataggatt 180
caaaagccct tcggatttct ggcagctctgt gcatagcacc ctgcctcggg agttggctcc 240
ttgtctagta tttaatacat ccccaaaactt ggctttattt tcagctgcct ttgccttcat 300
tgtgtgtaaa gatagtgcgg gtgactcaga tgtggtggtg caggagctca agtccatggt 360
ggccaccaag atcgccaaat atgctgtgcc tgatgagatc ctggtggtga aacgtcttcc 420
aaaaaccagg tctgggaagg tcatgcggcg gctcctgagg aagatcatca ctagtgaggc 480
ccaggagctg ggagacacta ccaccttgga ggacccagc atcatcgag agatcctgag 540
tgtctaccag aagtgcagg acaagcaggc tgctgctaag tgagctggca ccttggtggg 600
ctcttgggat gggcgggcac ccaagccctg gcttgctctt cccagaaggt acccctgagg 660
ttggcgtctt cctacgtccc agaagcagcc cccacccac acatgaccca caccgccctc 720
acgtgaagct gggctgagag ccctttctcc catccattgg aggtcccagg agtgtcaccc 780
atggagaggc tatgcgacat ggctagggct ggttctgcca tctgagtttg gtttcttgga 840
atgaaaaggc attgccatct ccattcctct gccctcttga gccagcacag gaagggtgagg 900
ccctgggata gcgcgcctgc tcagataaca cagagctagt tagctagtag caaccgtgtt 960
ttctccagat ctgtctagat acaaaggctca gaaatcttat ttttatactt ttatattgtg 1020
gaagaacagc atgcaacact cacatgtagt gtgtggattt acttgaacat gttcttttta 1080
acatgtagtt atgaaaatct ctttttttgc ctctactggt gaggaaacat gaggatcaga 1140
ggccacattt ttaattattg ttagtgtatt tggaaagtct aattggagat gtttgtacct 1200
ctgtctaaac agttcccttg agaacttcca agcctccggc atcttttctt ggtgagtggt 1260
tctcctgtgc ttggttgtgt ataattggagc taactcctaa gcggtggggg gaattgtggc 1320
gccttagttc tgaagctact ccagttatgt tctgtttctt caagctgtga tccagaaaga 1380
tttttgtgcc cccagatgcc tcttgatagg agaggcaaca tactccaaat agttgggttc 1440
ttcagggagc ctattagaaa ctcaggtgac ttgttagagc actaacttgg tcagagccaa 1500
atcctggcaa acgtgcctg accttcactc tctgggttgg gcagtgaaga ccactgagg 1560
ccaatgatga gacttggagg tctggatcca gtctctctt gttttaatgt gacttaggtg 1620
ctgtcaacat tagcaagata atggaaatca cgacgccagt ggggtgcttac ctccctgcta 1680
ggcatgcagg ggctggcggg tggcagggga aggagggcca gtgagccggg tcccttaggg 1740
gaggagagct ttgtcctctt tgcccacag tctacccttc agggccttgt ggcagtggca 1800
gtgttcgggg ggtgtctggg ccactgagta cccactcggg cgtggttggt ctggcctctt 1860
gggtgagtgaa acctgtgaag cccaggagggt ggtgttggt gcagggtaca caaatactga 1920
gtggtggtct tttgttacag gcttagcaac aaagctgtgc cctgggcatg gggggctgta 1980
gtgtagctac agttgtgctt ttgtgaaatg ccttagcttt ccattgtgct gagaggaacc 2040
tggacatggt cccgggcac tggatgatct gtaggggagg gagttcaaat aaagctttat 2100
tttgttcatt ttcaaaaaaa aa 2122

<210> 20
<211> 2357
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 2148623CB1

<400> 20
gctctttcag ggctgagcca tcttgcgtgt cttgcgctcg gtggaaatgc ccagccgagg 60
gacgcgacca gaggacagct ctgtgctgat cccacccgac aattcgaccc cacacaagga 120
ggatctaagc agcaagatta aagaacaaaa aattgtgggt gatgaacttt ctaaccttaa 180
gaagaatagg accgaacaga aatgctgtct gagagcaaga atatattgga tgaactgaaa 240
aaagaatacc aagaaataga aaacttagac aagacccaaa tcaagaaata gtcaacctga 300
tttcacataa caatgtgtgg catttgttgt tctgtaaact tttctgctga gcatttcagt 360
caagatttaa aagaggactt actatataat cttaaacagc ggggacccaa tagtagtaaa 420
caattgttaa agtctgatgt taactaccag tgtttatatt ctgctcacgt cctacacttg 480

```

aggggtgttt tgactaccca gcctgtggaa gatgaaagag gcaatgtgtt tctatggaat 540
ggagaaatTTT ttagtggaat aaaggttgaa gctgaagaga atgacactca aattttgttt 600
aattatcttt cctcctgtaa gaatgaatct gagattttgt cactcttctc agaagtacaa 660
ggccccgtgt catttatata ttatcaagca tctagtcatt atttatggtt tggtagggat 720
ttttttggtc gccgtagctt gctttggcat tttagtaatt tgggcaagag tttctgcctc 780
tcttcagttg gcacccaaac atctggattg gcaaatcagt ggcaagaagt tccagcatct 840
ggacttttca gaattgatct taagtctact gtcattttcca gatgcattat tttacaactg 900
tatccttgga aatatatttc tagggagaat attattgaag aaaaagttaa tagcctgagt 960
caaatttcag cagacttacc agcatttgta tcagtggtag caaatgaagc caaactgtat 1020
cttgaaaaac ctggtgttcc tttaaatatg atgttgccac aagctgcatt ggagactcat 1080
tgcagtaata tttccaatgt gccacctaca agagagatac ttcaagtctt tcttactgat 1140
gtacacatga aggaagtaat tcagcagttc attgatgtcc tgagtgtagc agtcaagaaa 1200
cgtgtcttgt gtttacctag ggatgaaaac ctgacagcaa atgaagtttt gaaaacgtgt 1260
gataggaaaag caaatgttgc aatcctgttt tctgggggca ttgattccat ggttattgca 1320
acccttgctg accgtcatat tcctttagat gaaccaattg atcttcttaa tgtagctttc 1380
atagctgaag aaaagaccat gccaaactacc tttaacagag aagggaataa acagaaaaat 1440
aaatgtgaaa taccttcaga agaattctct aaagatgttg ctgctgctgc tgctgacagt 1500
cctaataaac atgtcagtg accagatcga atcacaggaa gggcgaggact aaaggaacta 1560
caagctgtta gcccttcccg aatttggaa tttgttgaaa ttaatgtttc tatggaagaa 1620
ctgcagaaat taagaagaac tcgaatatgt cacttaattc ggccattgga tacagttttg 1680
gatgatagca ttggctgtgc agtctgtgtt gcttctagag gaattggttg gttagtggcc 1740
caggaaggag tgaataccta tcagagcaat gcaaaggtag ttctcactgg aattggtgca 1800
gatgagcaac tgcaggtta tctctgcat cgtgtccgct ttcagtcyca tgggctggaa 1860
ggattgaata aggaataaat gatggaactg ggtcgaattt cttctagaaa tcttggctgt 1920
gatgacagag ttattggtga tcatgaaaaa gaagcaagat ttcccttctc ggatgaaaaat 1980
gttgtctcct ttctaaattc tctgccgatt tgggaaaaag caaacttgac tttaccccca 2040
ggaattgggtg aaaaattact ttacgcctt ccagctgttg aacttggctc tacagcctct 2100
gctcttctgc ccaaacgggc catgcagttt ggatcaagaa ttgcaaaaat ggaaaaaatt 2160
aatgaaaagg catctgataa atgtggacgg ctccaaatca tgcctttaga aaatctttct 2220
attgaaaagg agactaaatt gtaatgtgat tcacaatgta acaatataaa aataagtttt 2280
tatataatta tataaaagta agatactctg ctgctttact attgtataat atagtagttt 2340
taaagttcaa aaaaaaa 2357

```

<210> 21

<211> 2136

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2579405CB1

<400> 21

```

ccgtccactt ggcgagtgag acgctgatgg gaggatggac atactggtgt ctgagtgtctc 60
cgcgcggtct ctgcagcagg aagaagagat taaatctctg actgctgaaa ttgaccgggtt 120
gaaaaactgt ggctgttttag gagcttctcc aaatttggag cagttacaag aagaaaaattt 180
aaaattaaag tatcgactga atattcttcg aaagagtcct caggcagaaa ggaacaaacc 240
aactaaaaat atgattaaca ttattagccg cctacaagag gtctttggtc atgcaattaa 300
ggctgcatac ccagatttgg aaaaatctcc tctgctagtg acaccaagtc agcaggccaa 360
gtttggggac tatcagtgtg atagtgtctat gggatattct cagatgtctc aaaccaagga 420
acagaaagtt aatccaagag aaattgtcga aaacattacc aaacacctcc cagacaatga 480
atgtattgaa aaagttgaaa ttgctgttcc tggttttatt aatgtccact taagaaagga 540
ttttgtatca gaacaattga ccagtcctct agtgaatgga gttcaactac ctgctctggg 600
agagaataaa aaggttatag ttgacttttc ctcccctaat atagctaaag agatgcatgt 660
aggccacctg aggtcaacta tcataggaga gagtataagc cgctcttttg aatttgcagg 720
gtatgacgtg ctcagggttaa atcatgtagg agactggggg acccagtttg gcatgtctat 780
cgctcacctg caagacaaat ttccagatta tctaacagtt tcacctccta ttggggatct 840
tcaggtcttt tataaggaat gtttgatact gaggaggaaat ttaagaagcg 900
agcatatcag tgtgtagttc tgctccaggg taaaaaccca gatattacaa aagcttggaa 960
gcttatctgt gatgtctccc gccaaagagt aaataaaatc tatgatgcat tggacgtctc 1020
tttaatatag agaggggaat ccttctatca agataggatg aatgatattg taaaggaatt 1080
tgaagataga ggatttgtgc aggtggatga tggcagaaag attgtatttg tcccagggtg 1140
ttccatacca ttaaccatag taaatcaga tggaggttat acctatgata catctgacct 1200
ggctgctatt aaacaaagac tatttgagga aaaagcagat atgattatct atgttgtgga 1260
caatggacaa tctgtgcact tccagacaa atttgtctgt gctcaaatga ttggttggtg 1320

```

```

tgaccctaaa gtaactcgag tcttccatgc tggatttggg gtggtgctag gggaagacaa 1380
gaaaaagttt aaaacacgtt cgggtgaaac agtgcgcctc atggatcttc tgggagaagg 1440
actaaaacga tccatggaca agttgaagga aaaagaaaga gacaagggtc taactgcaga 1500
ggaattgaat gctgctcaga catccgttgc atatggctgc atcaaatatg ctgaccttcc 1560
ccataaccgg ttgaatgact acatcttctc ctttgacaaa atgctagatg acagaggaaa 1620
tacagctgct tacttgttgt atgccttcac tagaatcagg tctattgcac gtctggccaa 1680
tattgatgaa gaaatgctcc aaaaagctgc tcgagaaacc aagattcttt tggatcatga 1740
gaaggaatgg aaactaggcc ggtgcatttt acggttccct gagattctgc aaaagatttt 1800
agatgactta tttctccaca ctctctgtga ttatatatat gagctggcaa ctgctttcac 1860
agagtcttat gatagctgct actgtgtgga gaaagataga cagactggaa aaatattgaa 1920
ggtgaacatg tggcgatgct tgctatgtga agcagtagct gctgtcatgg ccaaggggtt 1980
tgatatcctg ggaataaaac ctgtccaaag gatgtaatcc ttcataaggt tgaacactgt 2040
gtgtttttac caaagtggcc attggcactg tttgcttttt tacaatcatg tggacacaag 2100
cataagtaaa gaaaatttgt caaccaaaaa aaaaaa 2136

```

<210> 22

<211> 2480

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2662427CB1

<400> 22

```

ggaacgggtt tggcggtgtgg gacgcagctg cctctgtact ggggagtcac ggagtggccg 60
ggctccaggg acatggcggc ggctctctgc gtgtcgtgct tgcctggggc ggcggagagg 120
aacgggtggc atcgtctccc gagcctgctc gtgccgccga ggacatgggt gtggaggcaa 180
agaacatga agtacacaac agccacagga agaaacatta ccaaggctcct cattgcaaac 240
agaggagaaa ttgcctgcag ggtgatgcgc acagccaaaa aactgggtgt acagactgtg 300
tcggtttata gtgaggctga cagaaattcc atgcatgtag atatggcaga tgaagcatat 360
tccatcggcc ccgctccctc ccagcagagc tacctatcta tggagaaaat cattcaagt 420
gccaagacct ctgctgcaca ggctatccat ccaggatgcc gttttctttc agaaaacatg 480
gaatttgctg aactttgtaa gcaagaagga attattttta taggccctcc tccatctgca 540
attagagaca tgggtataaa gagcacatcc aaatccataa tggctgctgc tggagtacct 600
gttgtggagg gttatcatgg tgaggaccaa tcagaccagt gcctgaagga acacgccagg 660
agaattggct atcctgtcat gattaaagcc gtccgggggtg gaggaggaaa aggaatgagg 720
attgtagtat cagaacaaga atttcaagaa cagtttagat cagcacggag agaagctaag 780
aagtctttca atgatgatgc tatgctgata gagaagttg tagacacacc gaggcagtga 840
gaagtccagg tgtttgggtg tcaccatggc aatgctgtgt acttgtttga aagagactgt 900
agtgtgcaga ggcgacatca gaagatcatt gaggaggccc cagcgccctg tattaaatct 960
gaagtaagaa aaaagctggg agaagctgca gtcagagctg ctaaagctgt aaattatgtt 1020
ggagcaggga ctgtggagtt tattatggac tcaaaacata atttctgttt catggagatg 1080
aatacaaggc tgcaagtgga acatcctgtt actgagatga tcacagggaac tgacttggtg 1140
gagtggcagc ttagaattgc agcaggagag aagattcctt tgagccagga agaaataact 1200
ctgcagggcc atgccttcga agctagaata tatgcagaag atcctagcaa taacttcatg 1260
cctgtggcag gcccatagat gcacctctct actcctcgag cagacccttc caccaggatt 1320
gaaactggag tacggcaagg agacgaagtt tccgtgcatt atgaccccat gattgcgaag 1380
ctggtcgtgt gggcagcaga tcgccaggcg gcattgacaa aactgaggta cagccttcgt 1440
cagtacaata ttgttggact gccaccaaac attgacttct tactcaacct gtctggccac 1500
ccagagtttg aagctgggaa cgtgcacact gatttcatcc ctcaacacca caaacagttg 1560
ttgctcagtc ggaaggctgc agccaaagag tctttatgcc aggcagccct ggtctctatc 1620
ctcaaggaga aagccatgac cgacactttc actcttcagg cacatgatca attctctcca 1680
tttctgtcta gcagtggag aagactgaat atctcgtata ccagaaacat gactcttaaa 1740
gatggtaaaa acaatgtagc catagctgta acgtataacc atgatgggtc ttatagcatg 1800
cagattgaag ataaaaacttt ccaagtcctt ggtaatcttt acagcgaggg agactgcact 1860
tacctgaaat gttctgttaa tggagttgct agtaaagcga agctgattat cctggaaaac 1920
actatttacc tattttccaa ggaagggaagt attgagattg acattccagt ccccaaatac 1980
ttatcttctg tgagctcaca agaaactcag ggcggccctt tagctcctat gactggaacc 2040
attgaaaagg tgtttgtcaa agctggagac aaagtgaag cgggagattc cctcatggtt 2100
atgatcgcca tgaagatgga gcataccata aagtctccaa aggatggcac agtaaagaaa 2160
gtgttctaca gagaaggtgc tcaggccaac agacacactc ctttagtcga gtttgaggag 2220
gaagaatcag acaaaaaggga atcggaaata actccagcaa ggaaatggcc agttaagtag 2280
tgtcttctct ctccaccaa aagaggaagt gcctccagct tttctggggg tctcataaag 2340
agcagtttta ctaaatgatt gtatgcttat gctgaacacc tttcatattg gagaatcatg 2400

```

catttgggtc actaattatc tcaaaatatt tcataactaat aaagttgaat tattttttat 2460
 tggagccaa aaaaaaaaaa 2480

<210> 23
 <211> 2254
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2844928CB1

<400> 23
 ggccgggacg cagggcaaaag cgagccatgg ctgtctacgt cgggatgctg cgccctgggga 60
 ggctgtgcgc cgggagctcg ggggtgctgg gggccccggc cgccctctct cggagtggc 120
 aggaagccag gttgcagggt gtccgcttcc tcagttccag agaggtggat cgcattgtct 180
 ccacgcccc cggaggcctc agctacgttc aggggtgcac caaaaagcat cttaacagca 240
 agactgtggg ccagtgcctg gagaccacag cacagagggt ccagaacga gaggccttgg 300
 tcgtctcca tgaagacgtc aggttgacct ttgcccaact caaggaggag gtggacaaag 360
 ctgcttctgg cctcctgagc attggcctct gcaaaagtga ccggctgggc atgtggggac 420
 ctaactccta tgcattgggtg ctcatgcagt tggccaccgc ccaggcgggc atcattctgg 480
 tgtctgtgaa ccagcctac caggctatgg aactggagta tgtcctcaag aaggtgggct 540
 ggaagccctc ttgtttcccc aagcaattca agaccagca atactacaac gtcctgaagc 600
 agatctgtcc agaagtggag aatgcccagc caggggcctt gaagagtcag aggcctccag 660
 atctgaccac agtcattctc gtggatgccc ctttgccggg gaccctgtct ctggatgaag 720
 ttgtggcggc tggcagcaca cggcagcacc tggaccagct ccaatacaac cagcagttcc 780
 tgtcctgcca tgaccaccac aacatccagt tcacctcggg gacaacaggc agccccaagg 840
 ggccaccct ctccactac aacattgtca acaactccaa catttttaga gagcgcctga 900
 aactgcatga gaagacacca gagcagttgc ggatgacct gcccaacccc ctgtaccatt 960
 gcctgggttc cgtggcaggc acaatgatgt gtctgatgta cggtgccacc ctcatcctgg 1020
 cctctcccat cttcaatggc aagaaggcac tggaggccat cagcagagag agaggcacct 1080
 tcctgtatgg tacccccacg atgttcgtgg acattctgaa ccagccagac ttctccagtt 1140
 atgacatctc gaccatgtgt ggaggtgtca ttgctgggtc ccctgcacct ccagagtga 1200
 tccgagccat catcaacaag ataaatatga aggacctggt ggttgcttat ggaaccacag 1260
 agaacagtc cgtgacattc gcgcacttcc ctgaggacac tgtggagcag aaggcagaaa 1320
 gcgtgggcag aattatggc caccggagg cccggatcat gaacatggag gcagggacgc 1380
 tggcaaagct gaacacgccc ggggagctgt gcatccgagg gtactgcgtc atgctgggct 1440
 actgggggtga gcctcagaag acagaggaag cagtggatca ggacaagtgg tattggacag 1500
 gagatgtcgc cacaatgaat gagcagggtc tctgcaagat cgtgggccgc tctaaggata 1560
 tgatcatccg ggggtgtgag aacatctacc ccgagagct cgaggacttc ttccacacac 1620
 acccgaaggt gcaggaagtg caggtgaggc acttggtcct ggtgagcccc cagaaacaag 1680
 aaacacacat gaacacggtg atgtctgata tttttctttg gccctggaac gtgggtgggag 1740
 tgaaggacga tcggatgggg gaagagattt gtgctgcat tcggctgaag gacggggagg 1800
 agaccacggt ggagagata aaagctttct gcaaaaggaa gatctctcac ttcaagattc 1860
 cgaagtacat cgtgtttgtc acaaaactacc ccctcaccat ttcaggaaag atccagaaat 1920
 tcaaactctg agagcagatg gaacgacatc taaatctgtg aataaagcag caggcctgtc 1980
 ctggccggtt ggcttgactc tctcctgtca gaatgcaacc tggctttatg cacctagatg 2040
 tccccagcac ccagttctg gccaggcaca tcaaatgtca aggaattgac tgaacgaact 2100
 aagagctcct ggatgggtcc gggaactcgc ctgggcacaa ggtgccaaaa ggcaggcagc 2160
 ctgcccaggc cctccctcct gtccatcccc cacattcccc tgtctgtcct tgtgatttgg 2220
 cataaagagc ttctgttttc aaaaaaaaaa aaaa 2254

<210> 24
 <211> 1954
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 3231586CB1

<400> 24
 atccgggtacc acggccagtg caagctaaaa ttaaccctca ctaaaggga taagcttggc 60
 ccccgccgag atgtttcccc gcgagaagac gtggaacatc tcgttcgagg gctgcggctt 120
 cctcggegtc tactacgtcg gcgtggcctc ctgcctccgc gagcagcgc ccttctggtt 180

```

ggccaacgcc acgcacatct acggcgccctc ggccggggcg ctcacggcca cggcgctggt 240
caccgggggtc tgcctgggtg aggtggtgac caagttcatt gaggtatcta aagaggcccc 300
gaagcggttc ctggggcccc tgcacccctc cttcaacctg gtaaagatca tccgcagttt 360
cctgctgaag gtcctgcctg ctgatagcca tgagcatgcc agtgggcgcc tgggcatctc 420
cctgacccgc gtgtcagacg gcgagaatgt cattatatcc cacttcaact ccaaggacga 480
gtcatccag gccaatgtct gcagcggttt catccccgtg tactgtgggc tcatccctcc 540
ctccctccag ggggtgcgct acgtggatgg tggcatttca gacaacctgc cactctatga 600
gcttaagaac accatcacag tgtccccctt ctcgggcgag agtgacatct gtccgcagga 660
cagctccacc aacatccacg agctgcgggt caccaacacc agcatccagt tcaacctgcg 720
caacctctac cgctctccca agggcctctt cccgcgggag cccctggtgc tgcgagagat 780
gtgcaagcag ggataccggg atggcctgcg ctttctgcag cggaaacggc tcctgaaccg 840
gcccaacccc ttgctggcgt tgcacccccc cgcacccccc ggcccagagg acaaggacca 900
ggcagtgagg agcgcccaag cggaggatta ctcgcagctg cggggagaag atcacatcct 960
ggagcacctg cccgcccggc tcaatgagge cctgctggag gcctgcgtgg agcccacgga 1020
cctgctgacc accctctcca acatgctgcc tgtgcgtctg gccacggcca tgatggtgcc 1080
ctacacgctg ccgctggaga gcgctctgtc cttcaccatc cgcttgctgg agtggtgcc 1140
cgacgttccc gaggacatcc ggtggatgaa ggagcagacg ggcagcatct gccagtacct 1200
ggtgatgcgc gccaaagagg agctgggcag gcacctgccc tccaggctgc cggagcaggt 1260
ggagctgcgc cgcgtccagt cgtgcgcgtc cgtgcgcgtg tcctgcgcgc cctacagaga 1320
ggcactgccc ggctggatgc gcaacaacct ctcgctgggg gacgcgctgg ccaagtggga 1380
ggagtgcagg cgccagctgc tgcctggcct cttctgcacc aacgtggcct tcccgcccca 1440
agctctgcgc atgcgcgcac ccgcccagcc ggctcccgc ccccgggacc cagcatcccc 1500
gcagcacagg ctggccgggc ctagcccttt cctgagcacc cctgtctccc aggcccgccc 1560
cgtgatcggg gccctggggc tgtgagaccc cgacctctc gaggaacct gcctgagacg 1620
cctccattac cactgcgcag tgagatgagg ggactcacag ttgccaagag ggtctcttgc 1680
cgtgggcccc ctcgccagcc actcaccagc tgcattgact gagaggggag gtttccacac 1740
ccctcccctg ggccgctgag gccccgcgca cctgtgcctt aatcttccct ccctgtgct 1800
gcccgagcac ctccccgcc cctttactcc tgggaacttt gcagctgccc tcccccccc 1860
gtttttcatg gcctgctgaa atatgtgtgt gaagaattat ttattttcgc caaagcacat 1920
gtaataaatg ctgcagccca aaaaaaaaaa aaaa 1954

```

<210> 25

<211> 1937

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3580770CB1

<400> 25

```

gcagttcctg gggcgtagta ggggatccac aagcgtttgt gaccagtga gttctttaca 60
agggtgagat ctgcacggga ggacccgagc gaggggtctcg gcttgccagg aagccgggggt 120
tcccggggaa gcgtggagtt caccgcgca ctcgaaagtgc ctttgcaaaa ttatatctgg 180
gtgttggcac ccagccacta ttctgccaat gaagtacatc ctggtcacgg gtgggggtcat 240
ctcaggcatt ggtaaaggga tcattgccag cagcatttga acgattctaa aatcatgtgg 300
actccgagtt actgccataa aaatcgaccc ctatatatac atcgatgctg gcactttttc 360
accttatgaa cagggtgaag tcttcgtctt aaatgatggt ggagaagtgt atttagacct 420
tggagattat gaaagatttt tggatattaa tctttataaa gacacaatag tcaccacggg 480
gaagatatat cagcatgtga tcaataaaga gaggcgtggt gattacctgg ggaaaacagt 540
gcaagttgtc cctcacatta ctgatgctgt ccaggagtgg gttatgaatc aagccaaggt 600
gccggtggat ggtaataagg aagagcccca aatatgcgtt attgagctgg gaggcacat 660
tggagacatc gaaggaaatgc cgtttgtgga ggcgtttaga caattccagt ttaaggcgaa 720
aagagagaat ttctgtaata tccacgttag ccttgtccca cagctcagtg ctaccggaga 780
acaaaaaacc aaaccacccc aaaacagcgt ccgcgcactg aggggtttag gcctgtctcc 840
agatctgatt gtctgccgaa gttcaacgcc cattgagatg gccgtgaagg agaagatttc 900
tatgttttgt cagctgaacc ctgaacaggt catatgtatc catgatgttt ctccacata 960
ccgagttcct gtgcttttag aggaacaaaag cattgtgaaa tattttaagg agagattgca 1020
cctgcccacg ggtgatctct caagtaattt gctttttaag tggagaaata tggctgacag 1080
gtatgaaagg ttacagaaaa tatgctccat agccctggtt ggcaaataca ccaagctcag 1140
agactgctac gcctctgtgt tcaaagccct ggaacactca gccctggcca tcaaccacaa 1200
gttgaatctg atggtgatgg atatgcccga gcacaacctt ggcaatttgg gaggaacaat 1260
gagactggga ataagaagaa ctgttttcaa aactgaaaat tcaatattaa ggaaacttta 1320
tgggtgatgtt ccttttatag aagaaagaca cagacatcgg ttcgaggtaa accctaacct 1380
gatcaaacaa tttgagcaga atgacttaag ttttgtaggt caggatgttg atggagacag 1440

```



```

gatggaaatc attgaaactgg caaatcatcc ttattttgtt ggtgtccagt tccatcctga 1500
gttttcttct aggcgatga agccttcccc tccgtatctg gggctgttac ttgcagcaac 1560
tgggaacctg aatgcctact tgcaacaggg ttgcaaactg tcttccagt atagatacag 1620
tgatgccagt gatgacagct ttccagagcc aaggatagct gagttgaaa taagctgaaa 1680
tgaatacatg actgggaata atggggactg cctgtgaggg ctctgaaata attgaaggca 1740
agatgaagga actatctgaa gaaatcacta cactcttaga gaatccctct gttctccagc 1800
aaacatggga tgtaaacct caccagggaat ctgataatac atacttctgt caaccagaac 1860
cagaggggta gttttctttt ccctccagag gcaacctttg atacttaaaa tatctgtagc 1920
tgattaaatt tcgcca 1937

```

<210> 26

<211> 970

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3778612CB1

<400> 26

```

taataataat aacgtaatca tacctctagt catagcatac catttatcgg gctcggcgca 60
ggcccgcggg gagcgagcc cggcgagag actgatggag aggcagaaac ggaaggcgga 120
catcgagaaa gggctgcagt tcattcagtc gacactaccc cttaaagcaag aagaatatga 180
ggcctttctg ctcaagctgg tgcagaatct gtttgctgag ggcaatgatc tgttccggga 240
gaaggactat aagcaggtc tggtgcagta catggaaggg ctgaacgtgg ccgactacgc 300
tgctctgac caggtggccc tgccccggga gctgctgtgc aagctgcatg tcaatagggc 360
cgctgctac ttcaccatgg gcctgtatga gaaggcgctg gaggacagcg agaaggcgct 420
gggccctgac agtgagagta tccggcggtt gttccgcaa gcacgcgctc tcaatgaact 480
gggacgccac aaggagcgct acgagtgcag cagccggtgt tccctcgccc tgccccacga 540
tgaaagcggt actcagcttg gtcagggacc cttgggatct ggggcttctt ggccctggcca 600
gagctggagc cccacaggg taaggaagag agagtgggag gcagagtgtg atggggagga 660
gggacaggaa gacccttta atgatgagg taactatttc agttgtgagc cttctagggc 720
cccaggctgg gaggtcaga ggactgaatc tgggacctgt gttccccccg gcaggcaggg 780
acaagatggc atggcaagca tggggcggg gtgggtgggg agggatgctg catttctcag 840
ctgggcagta atcaatttaa tggtccttta aaatgtctgt gtattaaaaa tttaagaata 900
ccacacttta atattaaata ttcataaggt ctagtatctt gataataatg tagatgtttt 960
aataacaatt 970

```

<210> 27

<211> 1810

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4574912CB1

<220>

<221> unsure

<222> 193, 196-198

<223> a, t, c, g, or other

<400> 27

```

gctgcatgga gggcgccgtt ctagaagctg ggggtgcgcg gtgcttttgc aggtttgggt 60
gtgaattgag caagtatgaa aaccccggtt attcatcccc cagaagtgat tactttaaaa 120
attatatgat cattataact cagaatcgaa tgctcttct tgcaaatatg ttccacacga 180
tggactgtgt tgntgnnncc aggtatagct gtggaccaac tgtatatgat catgcgacc 240
ttggccatgc ttgctcatat gttagatttg atatcattcg aaggatccta accaaggttt 300
ttggatgcag catagtcatg gtgatgggta ttacagatgt agatgataaa atcatcaaaa 360
gagccaatga gatgaatatt tccccgctt cctcgcagc tctttatgag gaagacttca 420
agcaggacat ggcagccctg aaggttctcc caccacgggt gtacctgagg gtaaccgaaa 480
atattcctca gataattctt ttcattgaag gaatcattgc tcgtgggaac gcttattcaa 540
cggcaaaagg caatgtctac ttcgatctga agctagagg agacaagtat ggcaaatttg 600
tcggcggtgt cctgtgtcca gtcggagagc cagcggactc tgacaagcgt catgccagt 660
acttcgcctt gtggaaggcg gccaaaaccc aggaggtgtt ctgggcctct ccttggggac 720

```

```

ccgggaggcc gggctggcac atcgagtgt ctgccatcgc tagtatggt tttggaagtc 780
aactggatat ccattcaggt gggatagatt tagcttttcc acatcatgag aacgaaattg 840
cacagtgcga agtctttcat cagtgcgagc agtggggaaa ttattttctg cattctgggc 900
atttgacgcg caaaggcaaa gaagaaaaaa tgtccaaatc attaaagaac tacattacta 960
ttaaggactt tctgaagacc ttttcccccg atgtcttccg gttcttctgc ctgaggagca 1020
gctaccgctc agccatcgac tacagtgcga gcgccatgct ccaagctcag cagctgctcc 1080
tggggctggg ctcttttctg gaggacgcac gtgcctacat gaaggggcag ctggcctgag 1140
gctccgtcag ggaagcgatg ctgtgggaga ggctctccag caccaagagg gccgtgaagg 1200
gggccttggc agatgacttt gacacacca ggggtggtga tgccatcctg ggccttgac 1260
accaggggaa cggcagctc agggcgctcc tgaaggaaac tgaaggccg agaagtcctg 1320
ctgtgtttgg tgccatcatc tcttactttg aacagttttt tgaaactgtt ggaatttctc 1380
tggaacatca acagtacgtt tcaggagacg gcagcgaggc taccttgcat ggtgtggtgg 1440
acgagctggt gcggttccgg cagaaggtcc ggcagtttgc gctggccatg cccgaggcca 1500
cgggggacgc ccggcggcag cagctcctag aaaggcagcc cctgctggaa gcatgcgaca 1560
ccctgcgccc gggcctgact gccacggcga tcaacatcaa ggacagaagc agtacaacat 1620
ccacgtggga actgctggat caaaggacaa aagacaaaaa atcagcgggc tgaggatgga 1680
gcacagccat gaacctgtc acgacaagac gcacccatgc ttctcagggt caaggcttta 1740
tgtaaagct tcctgtcggg gctgtcaggt cagcattaaa gtaaggcaac caacagtga 1800
aaaaaaaaa

```

<210> 28

<211> 2162

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5630806CB1

<400> 28

```

gaccgtgcgc gcggcacgag gggacgggggt ccgactcaga aatggcgggc tccatgttct 60
acggcaggct agtggccgtg gccacccttc ggaaccaccg gcctcggaag gcccagcggg 120
ctgctgctca ggttctggga agttctggat tgtttaataa ccatggactc caagtacagc 180
agcaacagca aaggaatctc tcaactacatg aatacatgag tatggaatta ttgcaagaag 240
ctgggtgtctc cgttcccaaa ggatgtgtgg caaagtcacc agatgaagct tatgcaattg 300
ccaaaaaatt aggttcaaaa gatgtcgtga taaaggcaca ggttttagct ggtggttagg 360
gaaaaggaa atttgaaagt ggcctcaaag gaggagtga gatagtttct tctccagaag 420
aagcaaaagc tgtttcttca caaatgattg ggaaaaaatt gtttaccagg caaacgggag 480
aaaagggcag aatatgcaat caagtattgg tctgtgagcg aaaatatccc aggagagaat 540
actactttg aataacaatg gaaaggctat ttcaagggtc tgtattaata ggaagttcac 600
atggtgtgtg caacattgaa gatgttgcgt ctgagactcc tgaagcaata attaaagaac 660
ctattgatat tgaagaaggc atcaaaaagg aacaagctct ccagcttgca cagaagatgg 720
gatttccacc taatattgtg gaatcagcag cagaaaacat ggtcaagctt tacagccttt 780
ttctgaaata cgatgcaacc atgatagaaa taaatccaat ggtggaagat tcagatggag 840
ctgtattgtg tatggatgca aagatcaatt ttgactctaa ttcagcctat cgccaaaaga 900
aaatctttga tctacaggac tggacccagg aagatgaaag ggacaaagat gctgctaagg 960
caaatctcaa ctacattggc ctcgatggaa atataggctg cctagtaaat ggtgctgggt 1020
tggtatggc cacaatggat ataataaaac ttcattggagg gactccagcc aacttccttg 1080
atgttgggtg tgggtgtaca gtccatcaag taacagaagc atttaagctt atcacttcag 1140
ataaaaagg actggctatt ctgggtcaaca tttttggagg aatcatgcgc tgtgatgta 1200
ttgcacaggg tatagtcag gcagtaaaag acttggaaat taaaatacct gttgtgtgtac 1260
ggttacaagg tacacgagtc gatgatgcta aggcactgat agcggacagt ggacttaaaa 1320
tacttgcttg tgatgactg gatgaagctg ctagaatggt tgtaaagctc tctgaaatag 1380
tgaccttagc gaagcaagca catgtggatg tgaaatttca gttgccaata tgatctgaaa 1440
acccagtggg tggctgaagg tgttaaatgt gctataatca ttaagaatac tgtgttctgt 1500
gttattgttc tttttctttt tagtgtgtgg agattgtaat tgccatctag gcacacaaac 1560
atttaaaagg atttggactg catttaattg taccattcag aatggactgt ttgtacgaag 1620
catgtataat gcagttatct tctttcttcc gctcagccca gtcttttttg cttctcttac 1680
aaaacgtaac ttgcaatttg ccagtttatt attgttggat acaaagttct tcattgataa 1740
gagtcctata aataagataa atacgaagat aaagctttat tcttttagtg taaaatacag 1800
tatatcta atactagcctc attagtagag cagtatatata aaacaatgtt ttatgtaaaa 1860
agtgtttatc ttcagcacca aatacatgat aaatgtatca atcactatct ataaacagag 1920
ctttcaaaca ctctcagaa tattcttcta agtattttga tgaagtaact ttgtaattat 1980
ttgaacattg ttttaatcat taggaaacac tgattaactg caagtcttca tgattctgtc 2040
atattaagaa acacctgtag gtttgcttca aataaaggca tatataccaa ggacttacag 2100

```

acaaaattaa gaatgtcaat ttaagttaat aaaaatctcc caatatgaaa aaaaaaaaaa 2160
aa 2162

<210> 29
<211> 1477
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 5854855CB1

<400> 29
gcaatccgta cctcagtggt gtcccttttc agtgggttcc tttgtcccca ggcccattat 60
tccgtccctcc cctcttccct gatgtatttt ggcgcggtct cctgggtctg cgggcccagg 120
gctccggatg aggtctcccg ccgtcccgac ccccgcaagg gccagcttgg tgtcgcttc 180
gttcttctgc cacccttc ggaaggtgct cgggtcttcg gggcactggg tcccatcgg 240
cctcctcac ctggggtcac cctcggggtt ctggcctgta gcgagcaccg gctcagcaac 300
aagctgctgg ctggagcgg cgtcctcgag tggcaggaga agcgcagacc ctactctgac 360
tccactgcaa agctgaagcg gaccctgccc tgccaagcct acgtgaacca aggcgagaa 420
ctggagaccg accagtggcc gcagaagctg atcatgcagc tgatcccga gcagctgctg 480
accacctgg gcccctgtt ccggaactcc cagttggcac agttccactt caccaacaga 540
gactgcgact cgctcaagg gctctgccgc atcatgggca acggcttcgc gggctgcatg 600
ctgttcccc acatctcccc ctgtgaggtg ccgtgctca tgcctctgta ctgctccaag 660
aagaagatct tcatgggct catccctac gaccagagcg gcttcgtcag tggcatccg 720
caggtcatca ccaccgcaa gcagcgagtg ggacctggtg gtgtcaactc aggccagtc 780
cagatcgta acaacaagt tctggcatgg agtgggtgta tggagtggca ggagcccagg 840
cctgagccca acagtccgtc caagaggtgg ctgccatccc acgtctacgt gaaccagggg 900
gagatcctga ggaccgagca gtggccaagg aagctgtaca tgcagctcat cccgcagcag 960
ctgctgacca ccctagtgc gctgttcgg aactgcgcc tggctcagtt ccacttcacc 1020
aaggacctgg agacactgaa gagcctgtgc cggatcatgg acaatggctt cgccggctgc 1080
gtgcactttt cctacaaagc atcgtgtgag atccgcgtgc ttatgctcct gtactcttca 1140
gagaagaaaa tcttcattgg cctcatcccc catgaccagg gcaactttgt caacggcatc 1200
cgcggtgtca ttgccaacca gcagcaggtc ctgcagcggg acctggagca ggagcaacag 1260
caacgagggg tgggggggta gtggttacc cgggctgggc cctccagga gtacagatg 1320
agggccccgc agagactggt gacacgttc tgagcagggg cccctgggga cttcaactgc 1380
ccagcaacat ggaggatggt gtctgaggc ctccaaggac ggtccccacc cctctacgtt 1440
tcccaataa agccttttaa aaacctgaaa aaaaaaa 1477

<210> 30
<211> 1660
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 5993973CB1

<400> 30
gggccttccg cagctgcaga gcctcaacct cagcggcaac cggctgcgag agccgccagc 60
cgacctggcg cgctgcgccc cgcgcctgca gaccccaac ctcaccggca attgcctaga 120
ctcctttccc gccagctctt ttcgccccgg cgcgtgccc ctgctcagtg aactggcgcc 180
tgctgacaac tgctcccgag aactcagccc cgacatcgcc cacttgccct cgtcaagac 240
gttggaacct tcgaacaacc agctgagcga gatccctgca gagcttgagg actgccccaa 300
gctcaaggag atcaatttcc gtgggaacaa gctgagggac aagcgcctgg agaagatggt 360
cagcggctgc cagaccagat ccctcctgga gtacctgcgc gtcggaggcc gtggtggcgg 420
gaagggcaag ggcggtgccc agggctcgga gaaggaagag agccggagga agaggagga 480
gaggaagcag agggcggaag gtggtgatgg ggaggagcag gacgtgggag atgccggcgg 540
gctgctgctc agggctcctg acgtctctga aaacccgta cctctgacag tcagagttag 600
ccccgaggtc cgggatgtgc ggccctacat tgtggggggc gtggtgagag gcattggacct 660
gcagccaggg aatgcactca agcgttccct caccctgcag accaagctcc acgaagatct 720
ctgtgagaag aggaaggctg ccaccttgc caccacagag ctccgtgccc tcaaagggcc 780
cctgctgtac tgcgccccgc cccacagga cctcaagatt gtcccttgg ggcgaaaga 840
agccaaggcc aaggagctgg tgcggcagct gcagctggag gccgaggagc agaggaagca 900
gaagaagcgg cagagtgtgt cgggcctgca cagatacctt cacttgctgg atggaaatga 960

aaattaccg tgtcttgtg atgcagacg tgatgtgatt tccttccac caataacca 1020
cagtgagaag acaaaggta agaaaacgac ttctgatttg tttttggaag taacaagtgc 1080
caccagtctg cagatttgca aggatgtcat ggatgccctc attctgaaaa tggcagaaat 1140
gaaaaagtac actttagaaa ataaagagga aggatcactc tcagatactg aagccgatgc 1200
agtctctgga caacttccag atcccacaac gaatcccagt gctggaaagg acgggccctc 1260
ccttctgtg gtggagcagg tccgggtggt ggatctggaa gggagcctga aggtggtgta 1320
cccgtccaag gccgacctg ccactgcccc tccccacgtg actgtcgtgc gctgacgcca 1380
gggcccgtg tccgcgttg tttggccggt tttgcggagg tttctatgc gcaatgctga 1440
attatccgtt agattttcac ccagttttt ttgttggtt ttttttttg agatggagtc 1500
tcgtctgtc gccaggctg agtgcagtgg cgtgatctc agtcactgca gcctgtgtct 1560
cctgggttca agcgattctc ctgcctcagc ctcccaagta gctgggacta caggtgtgtg 1620
ccactaagct cagctaattt ttgtatttt agtagagacg 1660